

СИБИРСКИЕ ЭЛЕКТРОННЫЕ МАТЕМАТИЧЕСКИЕ ИЗВЕСТИЯ

Siberian Electronic Mathematical Reports

<http://semr.math.nsc.ru>

Том 15, стр. 768–772 (2018)

УДК 517.54

DOI 10.17377/semi.2018.15.062

MSC 30C65

ON COORDINATE VECTOR-FUNCTIONS OF QUASIREGULAR MAPPINGS

V.V. ASEEV

ABSTRACT. Let $f : R^n \rightarrow R^n = R^k \times R^{n-k}$ ($1 \leq k \leq n-1$) be a K -quasiregular mapping and $\pi : R^n \rightarrow R^k$ denotes the canonical projection. Then we obtain a lower estimate for the distortion of the values of generalized angles in R^k under the multy-valued function $F = f^{-1} \circ \pi^{-1} : R^k \rightarrow R^n$. This estimate is Möbius invariant and depends only on K and n .

Keywords: quasiregular map, conformal capacity of condenser, Teichmüller's ring, generalized angle, mapping of bounded angular distortion.

1. BASIC NOTIONS AND THE MAIN RESULT

The *generalized angle* in R^n is a four of sets $\Psi = (A_1, A_2; B_1, B_2)$ where $B_1 \neq \emptyset \neq B_2$, and $B_1 \cup B_2$ contains at least two distinct points. It's value is

$$\alpha(\Psi) := \inf_{a_1 \in A_1, a_2 \in A_2} \sup_{b_1 \in B_1, b_2 \in B_2} \frac{|a_1 - a_2| \cdot |b_1 - b_2|}{|a_1 - b_1| \cdot |a_2 - b_2| + |a_1 - b_2| \cdot |a_2 - b_1|},$$

where infimum through empty set is constituted to be equal 1. Note that $\alpha(\Psi) \leq 1$ for any generalized angle Ψ in R^n .

Set-valued mapping $F : X \rightarrow 2^Y$ is called *hyperinjective* if $x_1 \neq x_2$ is equivalent to $F(x_1) \cap F(x_2) = \emptyset$. The hyperinjective mapping $F : X \rightarrow 2^Y$ transforms generalized angle Ψ in X into generalized angle $F(\Psi)$ in Y , so in the case where $X \subset R^n$, $Y = R^n$ one can compare the values $\alpha(F(\Psi))$ and $\alpha(\Psi)$. Given an increasing real function $\omega : [0, 1] \rightarrow [0, 1]$ the hyperinjective mapping $F : X \rightarrow 2^Y$ is

ASEEV, V.V., ON COORDINATE VECTOR-FUNCTIONS OF QUASIREGULAR MAPPINGS.

© 2018 ASEEV V.V.

The work is supported by the program of fundamental scientific researches of the SB RAS No. 1.1.2., project No. 0314-2016-0007.

Received April, 17, 2018, published July, 16, 2018.

said to have ω -BAD property (Bounded Angular Distortion) if $\alpha(F(\Psi)) \geq \omega(\alpha(\Psi))$ for every generalized angle Ψ in X . The notion of generalized angle and its value was introduced in [1]. Some basic properties of generalized angles and mappings with BAD property in Ptolemaic spaces may be found in [1]-[2].

The definition and general properties of *quasiregular mappings* in R^n may be found, for example, in [3]-[5]. The advanced theory of quasiregular mappings was developed in well-known books [6]-[8], and for the information concerning with *conformal capacity of a condenser* and *the capacity of Teichmüller's extremal ring* the reader is referred to those monographs.

In order to formulate the main result we need the following:

Definition 1 (6, 3.2). *Compact proper subset $E \subset \bar{R}^n$ is said to be a set of null capacity, if for any compact set $B \subset \bar{R}^n \setminus E$ the conformal capacity $Cap(E, B)$ of the condenser (E, B) is equal to zero.*

Definition 2 (8, 7.18). *Given $e \in R^n$ with $|e| = 1$ the Teichmüller function $T_n(s)$ with parameter $s > 0$ is the conformal capacity of the condenser $(\{te : t \geq s\}, \{te : -1 \leq t \leq 0\})$ in R^n which is known as extremal Teichmüller's ring.*

This real function strictly decreases from $+\infty$ to 0 as s runs from 0 to $+\infty$.

Theorem 1 (4, 2.3). *Given a K -quasiregular mapping $f : D \rightarrow R^n$ of a domain $D \subset R^n$ with its complementary $\bar{R}^n \setminus D$ being a set of null capacity, for any open subset $A \subset D$ and a compact set $E \subset A$ the following inequality for conformal capacities holds*

$$Cap(f(E), \bar{R}^n \setminus f(A)) \leq K \cdot Cap(E, \bar{R}^n \setminus A) . \tag{1.1}$$

It follows that if $\bar{R}^n \setminus D$ is a set of null capacity then $\bar{R}^n \setminus f(D)$ is also a set of null capacity. Since a set of null capacity has zero topological dimension, the image $f(D) \subset R^n = R^k \times R^{n-k}$ under quasiregular mapping $f : D \rightarrow R^n$ mentioned in Theorem 1 has the canonical projection $\pi(f(D)) = R^k$ onto the whole coordinate k -plane R^k , $1 \leq k \leq n - 1$. So we can consider the set-valued hyperinjective mapping $F : R^k \rightarrow 2^D$, $F(x) = f^{-1}(\pi^{-1}(x))$, $x \in R^k$.

The $(n - 1)$ -dimensional Hausdorff measure of $\partial B(0, 1) \subset R^n$ will be denoted as Ω_{n-1}

Our main result is the following

Theorem 2. *Let a domain $D \subset R^n$ be such that its complement $\bar{R}^n \setminus D$ has null capacity and let $f : D \rightarrow R^n$ be a K -quasiregular mapping. Let $a \leq k \leq n - 1$ and $\pi : R^n = R^k \times R^{n-k} \rightarrow R^k$ be the canonical projection of R^n onto R^k . Then the set-valued mapping $F = f^{-1} \circ \pi^{-1} : R^k \rightarrow 2^D$ has the ω -BAD property with the control function*

$$\omega(t) = (3 - \sqrt{2}) \min \left\{ t, \exp \left[- \left(\frac{K \cdot \Omega_{n-1}}{T_n(1/t)} \right)^{\frac{1}{n-1}} \right] \right\} . \tag{1.2.1}$$

That is, the inequality

$$\alpha(F(\Psi)) \geq \omega(\alpha(\Psi)) \tag{1.2.2}$$

is valid for every generalized angle Ψ in R^k .

The similar result for the case $k = 1$ has been obtained in [2]. The counterexample in [2, §10] shows that in the case $k = n$ Theorem 2 is not valid with any control function ω that depends only on K and n .

2. AUXILIARY LEMMAS

Lemma 1. *Let a domain $D \subset \mathbb{R}^n$ be such that its complement $\bar{\mathbb{R}}^n \setminus D$ has null capacity. Let $\Phi = (\{z_1\}, \{z_2\}; P_1, P_2)$ be a generalized angle in D such that P_1 and P_2 are closed in D , z_1, z_2 are distinct points in $D \setminus (P_1 \cup P_2)$, and*

$$\alpha(\Phi) = t_0 < 3 - 2\sqrt{2}. \tag{2.1}$$

Then there exists a continuum $\gamma \subset D \setminus (P_1 \cup P_2)$ such that $z_1, z_2 \in \gamma$ and

$$\text{Cap}(\gamma, \bar{\mathbb{R}}^n \setminus (D \setminus (P_1 \cup P_2))) \leq \frac{\Omega_{n-1}}{[\log((1 - t_0)^2/(4t_0))]^{n-1}}. \tag{2.2}$$

Proof. The conformal capacity as well as the value of a generalized angle does not change under Möbius transformations in $\bar{\mathbb{R}}^n$. Thus we may assume without any loss of generality that $\infty \in P_1$, $z_1 = 0$, and $z_2 = e$ where $|e| = 1$. Thus the estimate

$$t_0 = \alpha(\Phi) = \sup_{b_1 \in P_1, b_2 \in P_2} \frac{|b_1 - b_2|}{|b_1| \cdot |b_2 - e| + |b_2| \cdot |b_1 - e|} \geq \frac{1}{|b| + |b - e|}$$

holds for all points $p \in P_2$. Then for every $b \in P_2$

$$2|b| + 1 \geq |b| + |b - e| \geq \frac{1}{t_0}; \quad |b| \geq \frac{1 - t_0}{t_0}$$

and it means that $P_2 \subset \{x \in \bar{\mathbb{R}}^n : |x| \geq (1 - t_0)/t_0\}$.

Choosing some point $q \in P_2$ we get the estimate

$$t_0 = \alpha(\Phi) \geq \frac{|b_1 - q|}{|q| \cdot |b_1 - e| + |b_1| \cdot |q - e|}$$

for all points $b_1 \in P_1$. Consequently

$$|q| - |b_1| \leq |b_1 - q| \leq t_0|q| \cdot |b_1 - e| + t_0|b_1| \cdot |q - e| \leq 2t_0|q| \cdot |b_1| + t_0|q| + t_0|b_1|$$

so that

$$|b_1| \geq \frac{(1 - t_0)|q|}{2t_0|q| + t_0 + 1} \geq \frac{(1 - t_0)^2}{4t_0}.$$

It means that $P_1 \subset \{x \in \bar{\mathbb{R}}^n : |x| \geq (1 - t_0)^2/(4t_0)\}$.

Thus $P_1 \cup P_2 \subset \{x : |x| \geq (1 - t_0)^2/(4t_0)\}$. The condition (2.1) gives the estimate $(1 - t_0)^2/(4t_0) > 1$, so that the sets $\bar{B}(0, 1)$ and $\bar{P}_1 \cup \bar{P}_2$ are separated by ring $\{x : 1 < |x| < (1 - t_0)^2/(4t_0)\}$.

Since $\bar{\mathbb{R}}^n \setminus D$ has zero topological dimension it does not break the connectivity of $\bar{B}(0, 1)$, so the points $z_1 = 0$ and $z_2 = e$ may be connected with a continuum $\gamma \subset D \cap \bar{B}(0, 1)$. The inequality

$$\begin{aligned} \text{Cap}(\gamma, \bar{P}_1 \cup \bar{P}_2) &\leq \text{Cap}(\gamma, (\bar{P}_1 \cup \bar{P}_2) \cup (\bar{\mathbb{R}}^n \setminus D)) \leq \\ &\text{Cap}(\gamma, \bar{P}_1 \cup \bar{P}_2) + \text{Cap}(\gamma, \bar{\mathbb{R}}^n \setminus D) = \text{Cap}(\gamma, \bar{P}_1 \cup \bar{P}_2) \end{aligned}$$

means that

$$\text{Cap}(\gamma, \bar{\mathbb{R}}^n \setminus (D \setminus (P_1 \cup P_2))) = \text{Cap}(\gamma, (\bar{\mathbb{R}}^n \setminus D) \cup (P_1 \cup P_2)) = \text{Cap}(\gamma, \bar{P}_1 \cup \bar{P}_2). \tag{2.3}$$

Since the ring $\{x : 1 < |x| < (1 - t_0)^2/(4t_0)\}$ separates γ from $\bar{P}_1 \cup \bar{P}_2$ the following estimate holds

$$\text{Cap}(\gamma, \bar{P}_1 \cup \bar{P}_2) \leq \frac{\Omega_{n-1}}{[\log((1 - t_0)^2/(4t_0))]^{n-1}}. \tag{2.4}$$

Now (2.3) and (2.4) give the desired estimate (2.2). □

Lemma 2. *Let $\Psi = (\{a_1\}, \{a_2\}; \{b_1\}, \{b_2\})$ be a generalized angle in R^k ($1 \leq k \leq n - 1$). In the euclidean space $R^n = R^k \times R^{n-k}$, consider the sets*

$$A_1 = \{a_1\} \times R^{n-k}; A_2 = \{a_2\} \times R^{n-k}; B_1 = \{b_1\} \times R^{n-k}; B_2 = \{b_2\} \times R^{n-k},$$

and a continuum $\Gamma \subset R^n \setminus (B_1 \cup B_2)$. If $\Gamma \cap A_1 \neq \emptyset \neq \Gamma \cap A_2$, then

$$\text{Cap}(\Gamma, B_1 \cup B_2) \geq T_n \left(\frac{1}{\alpha(\Psi)} \right) \tag{2.5}$$

Proof. Let $d = \min\{|a_1 - b_1|, |a_1 - b_2|\} = |a_1 - b_j|$ and $w_1 \in A_1 \cap \Gamma, w_2 \in A_2 \cap \Gamma$. There is a point $w_3 \in B_j$ with $|w_3 - w_1| = |a_1 - b_j| = d$. We use the well-known estimate (see [8, Lemma 7.34])

$$\text{Cap}(\Gamma, B_j) \geq T_n \left(\frac{|w_1 - w_3|}{|w_1 - w_2|} \right) \geq T_n \left(\frac{d}{|a_1 - a_2|} \right). \tag{2.6}$$

Since

$$\begin{aligned} \alpha(\Psi) &= \frac{|a_1 - a_2| \cdot |b_1 - b_2|}{|a_1 - b_1| \cdot |a_2 - b_2| + |a_1 - b_2| \cdot |a_2 - b_1|} \leq \\ &= \frac{|a_1 - a_2|}{d} \cdot \frac{|b_1 - b_2|}{|b_2 - a_2| + |a_2 - b_1|} \leq \frac{|a_1 - a_2|}{d}, \end{aligned}$$

we obtain from (2.6) the desired estimate (recall that T_n is a strictly decreasing function)

$$\text{Cap}(\Gamma, B_1 \cup B_2) \geq \text{Cap}(\Gamma, B_j) \geq T_n \left(\frac{d}{|a_1 - a_2|} \right) \geq T_n \left(\frac{1}{\alpha(\Psi)} \right).$$

□

3. PROOF OF THE MAIN THEOREM

It suffices to prove (1.2.2) for generalized angles $\Psi = (\{a_1\}, \{a_2\}; \{b_1\}, \{b_2\})$ in R^k (see [2, Lemma 4.2]). Consider the sets in $R^n = R^k \times R^{n-k}$

$$A_1 = \{a_1\} \times R^{n-k}; A_2 = \{a_2\} \times R^{n-k}; B_1 = \{b_1\} \times R^{n-k}; B_2 = \{b_2\} \times R^{n-k}.$$

Since $\bar{R}^n \setminus f(D)$ is a set of null capacity and has zero topological dimension, the sets $A'_j = A_j \cap f(D), B'_j = B_j \cap f(D)$ ($j = 1, 2$) are nonempty and closed in $f(D)$. Their preimages $Z_j = f^{-1}(A'_j) = F(a_j), P_j = f^{-1}(B'_j) = F(b_j)$ are nonempty mutually disjoint closed sets in D .

Since $\alpha(\Psi) \leq 1$ the inequality (1.2.2) holds in the case where $\alpha(F(\Phi)) \geq (3 - \sqrt{2})$. Thus we have to consider only the case where $\alpha(F(\Phi)) < (3 - \sqrt{2})$. In that case the inequality

$$\alpha(F(\Psi)) = \alpha(Z_1, Z_2; P_1, P_2) < 3 - \sqrt{2}$$

means that given any sufficiently small $\varepsilon > 0$ there exist points $z_1 \in Z_1, z_2 \in Z_2$ such that

$$t_0 := \alpha(\{z_1\}, \{z_2\}; P_1, P_2) < \alpha(F(\Psi)) + \varepsilon < 3 - \sqrt{2}.$$

Thus we have the situation described in Lemma 1 by which we have a continuum $\gamma \in D \setminus (P_1 \cup P_2)$ connecting z_1 to z_2 and satisfying the inequality (2.2). Since $B_1 \cup B_2 \subset \bar{R}^n \setminus f(D \setminus (P_1 \cup P_2))$ Theorem 1 may be applied to the open set $A = D \setminus (P_1 \cup P_2)$ and the compact set $E = \gamma$. Thus for the continuum $\Gamma = f(\gamma)$ we obtain the inequality

$$\text{Cap}(\Gamma, B_1 \cup B_2) \leq \text{Cap}(\Gamma, \bar{R}^n \setminus f(D \setminus (P_1 \cup P_2))) \leq K \cdot \text{Cap}(\gamma, \bar{R}^n \setminus (D \setminus (P_1 \cup P_2))).$$

Then Lemma 1 gives the estimate

$$\text{Cap}(\Gamma, B_1 \cup B_2) \leq \frac{K \cdot \Omega_{n-1}}{[\log((1-t_0)^2/(4t_0))]^{n-1}},$$

where $t_0 = \alpha(\{z_1\}, \{z_2\}; P_1, P_2)$. Now by Lemma 2 we have

$$T_n \left(\frac{1}{\alpha(\Psi)} \right) \leq \frac{K \cdot \Omega_{n-1}}{[\log((1-t_0)^2/(4t_0))]^{n-1}}. \quad (3.1)$$

Since $t_0 < 3 - \sqrt{2}$ it follows from (3.1) that

$$\frac{(2 - 2\sqrt{2})^2}{4t_0} \leq \exp \left[\left(\frac{K \cdot \Omega_{n-1}}{T_n(1/\alpha(\Psi))} \right)^{\frac{1}{n-1}} \right],$$

so that

$$\alpha(F(\Psi)) + \varepsilon > t_0 \geq (3 - 2\sqrt{2}) \exp \left[- \left(\frac{K \cdot \Omega_{n-1}}{T_n(1/\alpha(\Psi))} \right)^{\frac{1}{n-1}} \right]. \quad (3.2)$$

Since $\varepsilon > 0$ is arbitrarily close to 0 the desired estimate (1.2.1) follows immediately from (3.2). The Theorem 2 is proved.

The author is indebted to the referee for his/her valuable remarks, which have been used to improve the text of this paper.

REFERENCES

- [1] V.V. Aseev, A.V. Sychev, A.V. Tetenov, *Möbius-invariant metrics and generalized angles in Ptolemaic spaces*, Sib. Math. J., **46**:2 (2005), 189–204. MR2141193
- [2] V.V. Aseev, *Generalized angles in Ptolemaic Möbius structures*, Sib. Math. J., **59**:2 (2018), 241–256.
- [3] O. Martio, S. Rickman, J. Väisälä, *Definitions for quasiregular mappings*, Ann. Acad. Sci. Fenn. Ser. A I Math., **448** (1969), 1–40. MR0259114
- [4] O. Martio, S. Rickman, J. Väisälä, *Distortion and singularities of quasiregular mappings*, Ann. Acad. Sci. Fenn. Ser. A I Math., **465** (1970), 1–13. MR0267093
- [5] O. Martio, S. Rickman, J. Väisälä, *Topological and metric properties of quasiregular mappings*, Ann. Acad. Sci. Fenn. Ser. A I Math., **488** (1971), 1–31. MR0299782
- [6] Y.G. Reshetnyak, *Space mappings with bounded distortion*, Translations of Math. Monographs, **73** (1989), Amer. Math. Soc., Providence, R.I. MR0994644
- [7] S. Rickman, *Quasiregular mappings*, Springer-Verlag, Berlin, 1993. MR1238941
- [8] M. Vuorinen, *Conformal geometry and quasiregular mappings*, Lect. Notes Math., **1319** (1988), Springer-Verlag, Berlin-Heidelberg-New York-London-Paris-Tokio. MR0950174

VLADISLAV VASILYEVICH ASEEV
 SOBOLEV INSTITUTE OF MATHEMATICS,
 PR. KOPTYUGA, 4,
 630090, NOVOSIBIRSK, RUSSIA
E-mail address: `btp@math.nsc.ru`