

**CONSERVATIVE ALGEBRAS OF
2-DIMENSIONAL ALGEBRAS, V****IVAN KAYGORODOV, DOLORES MARTÍN BARQUERO,
AND CÁNDIDO MARTÍN GONZÁLEZ***Communicated by I.B. GORSHKOV*

Abstract: The notion of conservative algebras appeared in a paper of Kantor in 1972. Later, he defined the conservative algebra $W(n)$ of all algebras (i.e., bilinear maps) on the n -dimensional vector space. If $n > 1$, then the algebra $W(n)$ does not belong to any well-known class of algebras (such as associative, Lie, Jordan, or Leibniz algebras). It looks like that $W(n)$ in the theory of conservative algebras plays a similar role with the role of \mathfrak{gl}_n in the theory of Lie algebras. Namely, an arbitrary conservative algebra can be obtained from a universal algebra $W(n)$ for some $n \in \mathbb{N}$. The present paper is a part of a series of papers, which dedicated to the study of the algebra $W(2)$ and its principal subalgebras.

Keywords: bilinear maps, conservative algebra, graphs.

IVAN KAYGORODOV, DOLORES MARTIN BARQUERO, CÁNDIDO MARTIN GONZALEZ,
CONSERVATIVE ALGEBRAS OF 2-DIMENSIONAL ALGEBRAS, V.

© 2025 IVAN KAYGORODOV, DOLORES MARTIN BARQUERO, CÁNDIDO MARTIN GONZALEZ.

The work is supported by the Junta de Andalucía through projects UMA18-FEDERJA-119 and FQM-336 and by the Spanish Ministerio de Ciencia e Innovación through project PID 2019-104236GB-I00, all of them with FEDER funds; FCT 2023.08031.CEECIND, UIDB/MAT/00212/2020 and UIDP/MAT/00212/2020.

Received 18 Oct 2024, Published 16 June 2025.

Introduction

We will use the notation R for a (commutative) ring with unit. Also if M is an R -module and $n \geq 2$ an integer, we write $\text{Tor}_n(M) := \{x \in M : nx = 0\}$. The algebras under consideration in this work are not necessarily unital or associative. A multiplication on a vector space W is a bilinear mapping $W \times W \rightarrow W$. We denote by (W, P) the algebra with underlying space W and multiplication P . Given a vector space W , a linear mapping $A : W \rightarrow W$, and a bilinear mapping $B : W \times W \rightarrow W$, we can define a multiplication $[A, B] : W \times W \rightarrow W$ by the formula

$$[A, B](x, y) = A(B(x, y)) - B(A(x), y) - B(x, A(y)),$$

for $x, y \in W$. For an algebra A with a multiplication P and $x \in A$, we denote by L_x^P the operator of left multiplication by x . If the multiplication P is fixed, we write L_x instead of L_x^P .

In 1990 Kantor [15] defined the multiplication \cdot on the set of all algebras (i.e. all multiplications) on the n -dimensional vector space V_n as follows:

$$A \cdot B = [L_e^A, B],$$

where A and B are multiplications and $e \in V_n$ is some fixed vector. Let $W(n)$ denote the algebra of all algebra structures on V_n with multiplication defined above. If $n > 1$, then the algebra $W(n)$ does not belong to any well-known class of algebras (such as associative, Lie, Jordan, or Leibniz algebras). The algebra $W(n)$ turns out to be a conservative algebra (see below).

In 1972, Kantor [10] introduced conservative algebras as a generalization of Jordan algebras (also, see surveys about the study of conservative algebras and superalgebras [17, 26]). Namely, an algebra $A = (W, P)$ is called a *conservative algebra* if there is a new multiplication $F : W \times W \rightarrow W$ such that

$$[L_b^P, [L_a^P, P]] = -[L_{F(a,b)}^P, P],$$

for all $a, b \in W$. In other words, the following identity holds:

$$\begin{aligned} b(a(xy) - (ax)y - x(ay)) - a((bx)y) + (a(bx))y + (bx)(ay) - a(x(by)) \\ + (ax)(by) + x(a(by)) = -F(a, b)(xy) + (F(a, b)x)y + x(F(a, b)y). \end{aligned}$$

The algebra (W, F) is called an algebra *associated* to A . The main subclass of conservative algebras is the variety of terminal algebras, which is defined by the conservative identity with

$$F(a, b) = \frac{1}{3}(2ab + ba).$$

It includes the varieties of Leibniz and Jordan algebras as subvarieties.

Let us recall some well-known results about conservative algebras. In [10] Kantor classified all simple conservative algebras and triple systems of second-order and defined the class of terminal algebras as algebras satisfying a certain identity. He proved that every terminal algebra is a conservative algebra and classified all simple finite-dimensional terminal algebras with left quasi-unit over an algebraically closed field of characteristic zero [12]. Terminal trilinear operations were studied in [13], and some questions concerning the classification of simple conservative algebras were considered in [14]. After that, Cantarini and Kac classified simple finite-dimensional (and linearly compact) super-commutative and super-anticommutative conservative superalgebras and some generalizations of these algebras (also known as “rigid” or quasi-conservative superalgebras) over an algebraically closed field of characteristic zero [3]. The classification of all 2-dimensional conservative and rigid (in sense of Kac-Cantarini) algebras is given in [2]; and also, the algebraic and geometric classification of nilpotent low dimensional terminal algebras is given in [18, 19].

The algebra $W(n)$ plays a similar role in the theory of conservative algebras as the Lie algebra of all $n \times n$ matrices \mathfrak{gl}_n plays in the theory of Lie algebras. Namely, in [11, 15] Kantor considered the category \mathcal{S}_n whose objects are conservative algebras of non-Jacobi dimension n . It was proven that the algebra $W(n)$ is the universal attracting object in this category, i.e., for every $M \in \mathcal{S}_n$ there exists a canonical homomorphism from M into the algebra $W(n)$. In particular, all Jordan algebras of dimension n with unity are contained in the algebra $W(n)$. The same statement also holds for all noncommutative Jordan algebras of dimension n with unity. Some properties of the product in the algebra $W(n)$ were studied in [4, 16]. The universal conservative superalgebra was constructed in [21]. The study of low dimensional conservative algebras was started in [20]. The study of properties of 2-dimensional algebras is also one of popular topic in non-associative algebras (see, for example, [7, 23, 25]) and as we can see the study of properties of the algebra $W(2)$ could give some applications on the theory of 2-dimensional algebras. So, from the description of idempotents of the algebra $W(2)$ it was received an algebraic classification of all 2-dimensional algebras with left quasi-unit [22]. Derivations and subalgebras of codimension 1 of the algebra $W(2)$ and of its principal subalgebras W_2 and S_2 were described [20]. Later, the automorphisms, one-sided ideals, idempotents and local (and 2-local) derivations and automorphisms of $W(2)$ and its principal subalgebras were described in [1, 5, 22]. Note that W_2 and S_2 are simple terminal algebras with left quasi-unit from the classification of Kantor [12].


The present paper is devoted to continuing the study of properties of $W(2)$ and its principal subalgebras. We pay also some attention to the description of the affine group scheme of automorphisms of the algebras under scope, with an eye on the classification of gradings of these algebras (over arbitrary fields).

1 The graph of an algebra basis

In this section, the ground field \mathbb{F} will not be assumed to have characteristic zero. For an arbitrary \mathbb{F} -algebra A we will denote by $\mathcal{M}(A)$ (or simply \mathcal{M} if there is no possible ambiguity), the multiplication algebra of A , that is, the subalgebra of $\text{End}_{\mathbb{F}}(A)$ (where A is considered as a vector space) generated by left and right multiplication operators. We will denote by $\mathcal{M}_1(A)$ the subalgebra of $\text{End}_{\mathbb{F}}(A)$ generated by 1 and $\mathcal{M}(A)$. Observe that if A is an algebra whose multiplication algebra is \mathcal{M} , and $S \subset A$ a subset, the ideal of A generated by S agrees with $\mathcal{M}S$ (defined as the linear span of the elements $T(x)$ where $T \in \mathcal{M}$ and $x \in S$).

Assume A is an algebra over a field \mathbb{F} . Fix a basis $(u_i)_{i \in I}$ of A . Then we can construct a graph whose vertices are the basic elements u_i and for any two vertices we draw an arrow from u_i to u_j if $u_j = T(u_i)$ for some T in $\mathcal{M}_1(A)$. This relation $T(u_i) = u_j$ will be denoted $u_i \geq u_j$. The relation \geq is reflexive and transitive. However, to simplify the resulting graph, (1) we will not draw an arrow from each u_i to itself (as we should); and (2) if $u_i \geq u_j \geq u_k$ we will draw an arrow from u_i to u_j and another from u_j to u_k but there will be no need to draw the arrow from u_i to u_k . So there are many choices to draw the simplified graph, but they all give the same information.


For instance



is the graph associated to the algebra S_2 whose multiplication table (for a ground field of characteristic other than 3) is given below:

	e_1	e_2	e_3	e_4
e_1	$-e_1$	$-3e_2$	e_3	$3e_4$
e_2	$3e_2$	0	$2e_1$	e_3
e_3	$-2e_3$	$-e_1$	$-3e_4$	0
e_4	0	0	0	0

We can see that the graph is *strongly connected* in the sense that for any two vertices there is a path connecting them. This means that the ideal generated by any e_i is the whole algebra. In case the ground field has characteristic 3 the graph is given in the figure below,



Graph of S_2 in case $\text{ch}(\mathbb{F}) = 3$.

which is not strongly connected. As we will see, strong connection is a necessary condition for simplicity. If E is any graph with set of vertices E^0 and $S \subset E^0$, we will denote by $\mathbf{tree}(S)$ the set of all $v \in E^0$ such that there is a path from S to v . We can also construct a map $\mathbf{tree}: E^0 \rightarrow E^0$ such that $S \mapsto \mathbf{tree}(S)$. If E is the graph of an \mathbb{F} -algebra B relative to a basis $\mathcal{B} = \{b_i\}_i$, the fixed points of \mathbf{tree} induce ideals of B : assume $\mathbf{tree}(S) = S$, then $\oplus_{u \in S} \mathbb{F}u$ is a right ideal of B because for any $e_i \in \mathcal{B}$ and any $u \in S \subset \mathcal{B}$, one has $ue_i = 0$ or $ue_i = \sum_{j \in J} x^j e_j$ (with $x^j \in \mathbb{F}^\times$) so that $u \geq e_j$ (for any $j \in J$) implying $e_j \in S$. Thus $(\oplus_{u \in S} \mathbb{F}u)B \subset \oplus_{u \in S} \mathbb{F}u$. Similarly one can prove that $\oplus_{u \in S} \mathbb{F}u$ is a left ideal of B . So we claim:

Lemma 1. *Let E be the graph of an \mathbb{F} -algebra B relative to a given basis $\mathcal{B} = (b_i)_i$. If S is a fixed point of $\mathbf{tree}: E^0 \rightarrow E^0$ then $\oplus_{u \in S} \mathbb{F}u$ is an ideal of B .*

When $\text{ch}(\mathbb{F}) = 3$, considering the graph of S_2 in figure above, we see immediately that the unique fixed points of the map \mathbf{tree} are the subsets of vertices $\emptyset, \{e_1, e_3\}, \{e_1, e_2, e_3\}, \{e_1, e_3, e_4\}$ and E^0 . So at a first glance we detect three nontrivial proper ideals: $\mathbb{F}e_1 \oplus \mathbb{F}e_3, \mathbb{F}e_1 \oplus \mathbb{F}e_2 \oplus \mathbb{F}e_3$ and $\mathbb{F}e_1 \oplus \mathbb{F}e_3 \oplus \mathbb{F}e_4$. If I is the 3-dimensional ideal generated by e_1, e_3 and e_4 , we have $A = I \oplus \mathbb{F}e_2$ and A/I is a zero-product algebra. Similarly if J is the ideal generated by $\{e_1, e_2, e_3\}$ then $A = J \oplus \mathbb{F}e_4$ and A/J is a zero product algebra. With these ideas in mind, an easy criterium for simplicity is

Lemma 2. *Let A be any algebra with $A^2 \neq 0$ and let $\mathcal{M} := \mathcal{M}(A)$ be its multiplication algebra. Then A is simple if and only if*

- (1) *Its graph relative to a basis $(u_i)_{i \in I}$ is strongly connected, and*
- (2) *For any nonzero $x \in A$ there is some u_i in $\mathcal{M}x$.*

For instance to check the simplicity of the four-dimensional algebra S_2 whose multiplication table is given above (in the case of characteristic other than 3), since its graph is strongly connected we only need to realize that for a nonzero $x = \sum x_i e_i \in S_2$ we have:

- (1) If $x_3 \neq 0, (xe_2)e_2 = 3x_3e_2$ hence $e_2 \in \mathcal{M}(S_2)x$.
- (2) If $x_3 = 0, x_1 \neq 0, xe_2 = -3x_1e_2$, so $e_2 \in \mathcal{M}(S_2)x$.
- (3) If $x_3 = x_1 = 0, x_4 \neq 0, e_2x = x_4e_3$ implying $e_3 \in \mathcal{M}(S_2)x$.
- (4) If $x_i = 0$ except for $i = 2$ then $e_2 \in \mathcal{M}(S_2)x$.

Thus, in any case there is a basis element in $\mathcal{M}(S_2)x$.

For an algebra A , the condition that $\mathcal{M}(A) = \text{End}_{\mathbb{F}}(A)$ implies simplicity of A : indeed, if this coincidence happens, for any nonzero $x \in A$ and any $y \in A$, there is a linear map $f: A \rightarrow A$ such that $y = f(x)$. Since $f \in \mathcal{M}(A)$ then y is in the ideal generated by x . Thus A is simple. In [8, Corollary of Theorem 3] it is proved that in the finite-dimensional case, an algebra U over a field \mathbb{F} is simple if and only if its multiplication algebra is simple.

Lemma 3. *Let U be a finite-dimensional algebra over a field \mathbb{F} . If U is simple then $\mathcal{M}(U) = \text{End}_{\mathbb{F}}(U)$.*

Proof. Assume first that the ground field \mathbb{F} is algebraically closed. If U is simple, by [8] we know that $\mathcal{M} := \mathcal{M}(U)$ is simple. Let $n := \dim(U)$, since U is an \mathcal{M} -module (irreducible and faithful) then $\mathcal{M} \cong \text{End}_{\mathbb{F}}(U)$. If the ground field \mathbb{F} is not algebraically closed we consider the algebraic closure Ω of \mathbb{F} and the Ω -algebra $U_{\Omega} := U \otimes_{\mathbb{F}} \Omega$. Then $\mathcal{M}(U_{\Omega}) = \text{End}_{\Omega}(U_{\Omega})$ and by [6, (2.5)Lemma] we have $\mathcal{M}(U_{\Omega}) \cong \mathcal{M}(U) \otimes \Omega$. Since

$$\dim(\text{End}_{\mathbb{F}}(U)) = \dim_{\Omega}(\text{End}_{\Omega}(U_{\Omega})) = \dim_{\Omega}(\mathcal{M}(U_{\Omega})) = \dim(\mathcal{M}(U))$$

we conclude $\text{End}_{\mathbb{F}}(U) = \mathcal{M}(U)$. □

If U is simple but fails to be finite-dimensional we can say that $\mathcal{M}(U)$ is a dense subalgebra of $\text{End}_{\mathbb{F}}(U)$ in the sense of Jacobson density. To clarify this, the \mathcal{M} -module U is simple (or irreducible in the terminology of [9]). Since the action of \mathcal{M} on U is the natural one, we can say that U is an irreducible and faithful \mathcal{M} -module. Hence \mathcal{M} is a primitive \mathbb{F} -algebra. The irreducibility of U as an \mathcal{M} -module implies that the \mathbb{F} -algebra $\Gamma := \text{End}_{\mathcal{M}}(U)$ is a division algebra. This consists of all \mathbb{F} -linear maps $T: U \rightarrow U$ such that

$$T(xy) = xT(y) = T(x)y \text{ for any } x, y \in U.$$

So Γ is the centroid of U which is known to be a field (extension of \mathbb{F}) given the simplicity of U . Now U is an Γ -algebra in a natural way and we have a monomorphism $\mathcal{M} \hookrightarrow \text{End}_{\Gamma}(U)$ which is dense in the sense that for any Γ -linearly independent $x_1, \dots, x_n \in U$ and arbitrary $y_1, \dots, y_n \in U$, there is some $T \in \mathcal{M}$ such that $T(x_i) = y_i$ for $i = 1, \dots, n$ (see [9, Density Theorem for Irreducible Modules, II, §2, p.28]). Observe that when U is finite-dimensional the extension field Γ has $(\Gamma : \mathbb{F})$ finite. Thus if \mathbb{F} is algebraically closed we have $\Gamma = \mathbb{F}$ and \mathcal{M} being dense in $\text{End}_{\Gamma}(U) = \text{End}_{\mathbb{F}}(U)$ gives $\mathcal{M} = \text{End}_{\mathbb{F}}(U)$. If \mathbb{F} is not algebraically closed we can argue as in the last part of the proof of Lemma 3. So we recover Lemma 3 from the general result:

Proposition 4. *If U is a simple \mathbb{F} -algebra then $\mathcal{M} := \mathcal{M}(U)$ is a primitive algebra, more precisely there is:*

- (1) *A monomorphism $\mathcal{M} \hookrightarrow \text{End}_{\Gamma}(U)$ where Γ is the centroid of U (which is a field extension of \mathbb{F}).*
- (2) *For any collection of Γ -linearly independent elements $x_1, \dots, x_n \in U$ and any collection $y_1, \dots, y_n \in U$, there is an element $T \in \mathcal{M}$ such that $T(x_i) = y_i$ for any i .*

As a consequence of Lemma 3, for a finite-dimensional algebra A over a field \mathbb{F} , proving that $\text{End}_{\mathbb{F}}(A)$ agrees with $\mathcal{M}(A)$ is equivalent to proving that A is simple. The characterization of the coincidence $\mathcal{M}(A) = \text{End}_{\mathbb{F}}(A)$ in terms of the graph of A is:

Proposition 5. *Let $A^2 \neq 0$ be a finite-dimensional algebra and $\mathcal{M} = \mathcal{M}(A)$ its multiplication algebra. Then $\mathcal{M} = \text{End}_{\mathbb{F}}(A)$ if and only if:*

- (1) *The graph of A relative to a basis $(u_i)_{i=1}^n$ is strongly connected.*

- (2) For every $i \in \{1, \dots, n\}$ there is some $j \in \{1, \dots, n\}$ and $T \in \mathcal{M}$ such that $T(u_k) = \delta_{ik}u_j$ for any k .

Proof. If $\mathcal{M} = \text{End}_{\mathbb{F}}(A)$ the algebra A is simple whence the graph is strongly connected. The other assertion in the statement is straightforward. So assume that both conditions in the statement hold. If we define the linear maps $E_{ij}: A \rightarrow A$ such that $E_{ij}(u_k) = \delta_{ik}u_j$ we know that $\text{End}_{\mathbb{F}}(A) = \bigoplus_{i,j=1}^n \mathbb{F}E_{ij}$. Now, condition 2) says that for any i there is some $E_{ij} \in \mathcal{M}$. But the graph relative to (u_i) is strongly connected so for any u_j, u_k there exists $T \in \mathcal{M}$ such that $u_k = T(u_j)$. Thus $TE_{ij} = E_{ik}$ and we have $E_{ik} \in \mathcal{M}$ for every k and i . \square

Remark 6. If the graph of an \mathbb{F} -algebra A relative to a basis $(u_i)_{i=1}^n$ is strongly connected and some $E_{ij} \in \mathcal{M} = \mathcal{M}(A)$ (identifying \mathcal{M} with an subalgebra of $\text{End}_{\mathbb{F}}(A)$), then $E_{ik} \in \mathcal{M}$ for any k . Indeed: given u_j and u_k by the strong connectedness of the graph, there is some $T \in \mathcal{M}$ such that $T(u_j) = u_k$. Then $E_{ik} = TE_{ij} \in \mathcal{M}$.

Theorem 7. If \mathbb{F} is a field of characteristic other than 3 and S_2 the four-dimensional algebra whose multiplication algebra is given above, we have $\mathcal{M} := \mathcal{M}(S_2) = \text{End}_{\mathbb{F}}(S_2)$. Consequently S_2 is simple. If the characteristic of \mathbb{F} is 3 there is a 3-dimensional ideal I which is the subspace generated by e_1, e_3 and e_4 . Moreover $S_2 = I \oplus \mathbb{F}e_2$ and S_2/I is a zero-product algebra. In this case \mathcal{M} has dimension 8 and a 4-dimensional radical $\text{rad}(\mathcal{M})$ such that $\text{rad}(\mathcal{M}) = 0$ and $\mathcal{M}/\text{rad}(\mathcal{M}) \cong M_2(\mathbb{F})$.

Proof. Since the graph relative to the basis $(e_i)_{i=1}^4$ is strongly connected we need to check (2) in Proposition 5.

- (A) First, we will consider the case in which the characteristic of \mathbb{F} is other than 2 or 3. Under this hypothesis, the element $R_{e_2}^2 \in \mathcal{M}$ acts in the way

$$\begin{aligned} R_{e_2}^2(e_1) &= (e_1e_2)e_2 = -3e_2^2 = 0, & R_{e_2}^2(e_2) &= (e_2e_2)e_2 = 0, \\ R_{e_2}^2(e_3) &= (e_3e_2)e_2 = -e_1e_2 = 3e_2, & R_{e_2}^2(e_4) &= (e_4e_2)e_2 = 0. \end{aligned}$$

Thus $E_{32} = \frac{1}{3}R_{e_2}^2 \in \mathcal{M}$ and we can also prove that $E_{3k} \in \mathcal{M}$ for any k :

$$E_{31} = \frac{1}{2}R_{e_3}E_{32} \in \mathcal{M}, \quad E_{33} = R_{e_4}E_{32} \in \mathcal{M}, \quad E_{34} = -\frac{1}{3}R_{e_3}E_{33} \in \mathcal{M}.$$

So far $E_{3k} \in \mathcal{M}$ for any k . Furthermore, it can be checked that

$$R_{e_1} = -E_{11} + 3E_{22} - 2E_{33}, \quad L_{e_1} = -E_{11} - 3E_{22} + E_{33} + 3E_{44}.$$

On the other hand we have:

$$\begin{aligned} E_{21} &= -R_{e_2}R_{e_4} \in \mathcal{M}, & & \\ R_{e_2} &= -3E_{12} - E_{31}, & & \text{hence } E_{12} \in \mathcal{M} \\ E_{12}, E_{21} &\in \mathcal{M}, & & \text{hence } E_{11}, E_{22} \in \mathcal{M} \end{aligned}$$

$$L_{e_1} = -E_{11} - 3E_{22} + E_{33} + 3E_{44}, \quad \text{hence } E_{44} \in \mathcal{M}.$$

Summarizing $E_{ii} \in \mathcal{M}$ for $i = 1, 2, 3, 4$. Thus (2) of Proposition 5 is satisfied.

(B) Second, we analyze the case in which \mathbb{F} has characteristic 2. The multiplication table of S_2 adopts the form in figure:

	e_1	e_2	e_3	e_4
e_1	e_1	e_2	e_3	e_4
e_2	e_2	0	0	e_3
e_3	0	e_1	e_4	0
e_4	0	0	0	0

Multiplication table of S_2 when $\text{char}(\mathbb{F}) = 2$.

Then the matrix whose (i, j) entry is $R_{e_i}R_{e_j}$ is shown in

$$(R_{e_i}R_{e_j})_{i,j=1}^4 = \begin{pmatrix} E_{11} + E_{22} & E_{12} + E_{31} & 0 & 0 \\ E_{12} & E_{32} & E_{11} & E_{21} \\ E_{13} & E_{33} & E_{14} & E_{24} \\ E_{14} + E_{23} & E_{13} + E_{34} & 0 & 0 \end{pmatrix},$$

hence $E_{1i} \in \mathcal{M}$ for any i . From this, $E_{2i} \in \mathcal{M}$ also for any i . Consequently $E_{3i} \in \mathcal{M}$ for any i . The matrix whose (i, j) entry is $L_{e_i}L_{e_j}$ is

$$(L_{e_i}L_{e_j})_{i,j=1}^4 = \begin{pmatrix} E_{11} + E_{22} + E_{33} + E_{44} & E_{12} + E_{43} & E_{21} + E_{34} & 0 \\ E_{12} + E_{43} & 0 & E_{22} + E_{33} & 0 \\ E_{21} + E_{34} & E_{11} + E_{44} & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix},$$

which implies $E_{43}, E_{44} \in \mathcal{M}$. $E_{41}, E_{42} \in \mathcal{M}$ follows from the multiplication table.

(C) Third, assume that $\text{char}(\mathbb{F}) = 3$. Denoting as before by E_{ij} the basis of $\text{End}_{\mathbb{F}}(A)$ such that $E_{ij}(e_k) = \delta_{ik}e_j$, we have

$$\begin{aligned} L_{e_1} = R_{e_1} &= -E_{11} + E_{33}, & L_{e_2} &= -E_{31} + E_{43}, & R_{e_2} &= -E_{31}, \\ L_{e_3} = R_{e_3} &= E_{13} - E_{21}, & R_{e_4} &= E_{23}. \end{aligned}$$

The subalgebra of $\text{End}_{\mathbb{F}}(A)$ generated by these operators is \mathcal{M} and coincides with the \mathbb{F} -linear span of

$$\{E_{11}, E_{33}, E_{31}, E_{13}, E_{21}, E_{23}, E_{41}, E_{43}\}.$$

If we compute the radical of the symmetric bilinear form $\langle \cdot, \cdot \rangle: \mathcal{M} \times \mathcal{M} \rightarrow \mathbb{F}$ given by $\langle f, g \rangle := \text{trace}(fg)$ we find that

$$\mathcal{M}^\perp := \text{rad}(\langle \cdot, \cdot \rangle) = \mathbb{F}E_{43} \oplus \mathbb{F}E_{21} \oplus \mathbb{F}E_{23} \oplus \mathbb{F}E_{41}.$$

For the reader's convenience, we recall that the radical of a bilinear symmetric form in a vector space is the subspace of elements which

are orthogonal to the whole space. For a finite-dimensional associative algebra of endomorphisms, the bilinear form $\langle f, g \rangle := \text{trace}(fg)$ is associative in the sense that $\langle fg, h \rangle = \langle f, gh \rangle$. Then, it is easy to realize that \mathcal{M}^\perp is an ideal of the algebra. Now, one can see that

$$(\mathcal{M}^\perp)^2 = 0 \text{ and } \mathcal{M}/\mathcal{M}^\perp \cong \mathbb{F}E_{11} \oplus \mathbb{F}E_{13} \oplus \mathbb{F}E_{31} \oplus \mathbb{F}E_{33} \cong M_2(\mathbb{F}).$$

Thus \mathcal{M}^\perp is a maximal ideal and since it is nilpotent we conclude that \mathcal{M}^\perp is the radical of \mathcal{M} . □

2 Automorphisms and multiplication algebras

2.1. Conservative algebra S_2 . In this subsection, we compute the automorphism group scheme of S_2 over a field \mathbb{F} of arbitrary characteristic. In order to do that we consider an associative, commutative and unital ring R and define in the free R -module B with basis $\{e_i\}_{i=1}^4$ the product as in table of multiplication of S_2 (extended by R -bilinearity to the whole B). When $R = \mathbb{F}$ the algebra B is precisely S_2 . So by considering B we are thinking about A defined over an arbitrary ring R (associative, commutative and unital). If we are able to determine $\text{aut}_R(B)$, then we have walked a long way towards the knowledge of the affine group scheme of A over arbitrary fields. So consider $f \in \text{aut}(B)$, and write $f(e_i) = f_i^j e_j$ (using Einstein Criterion). Then $f(e_1^2) = f(e_1)^2$, hence

$$\begin{aligned} -f(e_1) &= (f_1^j e_j)^2 = -(f_1^1)^2 e_1 - 3(f_1^3)^2 e_4 - 3f_1^1 f_1^2 e_2 + f_1^1 f_1^3 e_3 + 3f_1^1 f_1^4 e_4 + \\ &\quad 3f_1^2 f_1^1 e_2 + 2f_1^2 f_1^3 e_1 + f_1^2 f_1^4 e_3 - 2f_1^3 f_1^1 e_3 - f_1^3 f_1^2 e_1 = \\ &\quad [-(f_1^1)^2 + f_1^2 f_1^3] e_1 + [f_1^2 f_1^4 - f_1^1 f_1^3] e_3 + [-3(f_1^3)^2 + 3f_1^1 f_1^4] e_4, \end{aligned}$$

so we deduce

$$\begin{cases} f_1^1 = (f_1^1)^2, & f_1^2 = 0 \\ f_1^3 = f_1^1 f_1^3, & f_1^4 = 3(f_1^3)^2 - 3f_1^1 f_1^4 \end{cases}$$

Furthermore, since Re_4 is the left annihilator of B (and this is preserved under automorphism), we have $f(e_4) = f_4^4 e_4$, which implies $f_4^4 \in R^\times$ (invertible elements of R). Now applying f to $e_1 e_4 = 3e_4$ we get

$$(f_1^1 e_1 + f_1^3 e_3 + f_1^4 e_4) e_4 = 3e_4, \text{ that is, } 3f_1^1 e_4 = 3e_4.$$

Assume now that $\text{Tor}_3(R) = 0$, then $f_1^1 = 1$ so that $4f_1^4 = 3(f_1^3)^2$.

Then we discuss cases:

- (A) $\frac{1}{3}, \frac{1}{2} \in R$. Then $f_1^4 = \frac{3}{4}(f_1^3)^2$ (besides $f_1^1 = 1, f_1^2 = 0$). Since $e_3 e_4 = 0$ we have $f(e_3) e_4 = 0$ hence $f_3^1 e_1 e_4 + f_3^2 e_2 e_4 = 0$. So $3f_3^1 e_4 + f_3^2 e_3 = 0$ implying $f_3^1 = f_3^2 = 0$. So far the matrix of f relative to the R -basis $\{e_i\}$ is

$$\begin{pmatrix} 1 & 0 & * & * \\ f_2^1 & f_2^2 & f_2^3 & f_2^4 \\ 0 & 0 & f_3^3 & * \\ 0 & 0 & 0 & f_4^4 \end{pmatrix} \tag{1}$$

whose determinant is $f_2^2 f_3^3 f_4^4$ and must be in R^\times . Hence $f_i^i \in R^\times$ for any i . Also $e_3^2 = -3e_4$ and applying f we have

$$(f_3^3 e_3 + f_3^4 e_4)^2 = -3f_4^4 e_4.$$

Thus $-3(f_3^3)^2 e_4 = -3f_4^4 e_4$ which gives $f_4^4 = (f_3^3)^2$. Take now into account that $e_3 e_2 = -e_1$ hence

$$(f_3^3 e_3 + f_3^4 e_4)(f_2^i e_i) = -(e_1 + f_1^3 e_3 + f_1^4 e_4)$$

or equivalently $f_3^3 e_3 (f_2^i e_i) = -(e_1 + f_1^3 e_3 + f_1^4 e_4)$. So

$$-2f_3^3 f_2^1 e_3 - f_3^3 f_2^2 e_1 - 3f_3^3 f_2^3 e_4 = -e_1 - f_1^3 e_3 - f_1^4 e_4$$

and we get

$$f_2^2 f_3^3 = 1, \quad f_1^3 = 2f_3^3 f_2^1, \quad f_1^4 = 3f_3^3 f_2^3.$$

Now $e_2^2 = 0$ hence $(f_2^i e_i)^2 = 0$. Thus

$$\begin{aligned} &-(f_2^1)^2 e_1 - 3(f_2^3)^2 e_4 - 3f_2^1 f_2^2 e_2 + f_2^1 f_2^3 e_3 + 3f_2^1 f_2^4 e_4 + \\ &3f_2^2 f_2^1 e_2 + 2f_2^2 f_2^3 e_1 + f_2^2 f_2^4 e_3 - 2f_2^3 f_2^1 e_3 - f_2^3 f_2^2 e_1 = 0. \end{aligned}$$

We get

$$-(f_2^1)^2 + f_2^2 f_2^3 = 0, \quad f_2^2 f_2^4 - f_2^3 f_2^1 = 0, \quad -3(f_2^3)^2 + 3f_2^1 f_2^4 = 0.$$

Also $e_1 e_3 = e_3$ so that $(e_1 + f_1^3 e_3 + f_1^4 e_4) f_3^i e_i = f_3^i e_i$. Equivalently

$$(e_1 + f_1^3 e_3)(f_3^3 e_3 + f_3^4 e_4) = f_3^3 e_3 + f_3^4 e_4.$$

Then $f_3^3 e_3 + 3f_3^4 e_4 - 3f_1^3 f_3^3 e_4 = f_3^3 e_3 + f_3^4 e_4$ and we get $2f_3^4 - 3f_1^3 f_3^3 = 0$ so that $f_3^4 = \frac{3}{2} f_1^3 f_3^3$. Thus if we put $f_2^2 = \lambda$ and $f_1^3 = \mu$ we have $f_3^3 = \frac{1}{\lambda}$ and

$$\begin{aligned} f_1^4 &= \frac{3}{4} \mu^2, & f_2^1 &= \frac{\lambda \mu}{2}, & f_2^3 &= \frac{\lambda \mu^2}{4}, \\ f_2^4 &= \frac{\lambda \mu^3}{8}, & f_3^1 &= f_3^2 = 0, & f_3^4 &= \frac{3\mu}{2\lambda}. \end{aligned}$$

Thus the matrix of f in the basis $\{e_i\}$ is

$$w(\lambda, \mu) = \begin{pmatrix} 1 & 0 & \mu & \frac{3\mu^2}{4} \\ \frac{\lambda \mu}{2} & \lambda & \frac{\lambda \mu^2}{4} & \frac{\lambda \mu^3}{8} \\ 0 & 0 & \frac{1}{\lambda} & \frac{3\mu}{2\lambda} \\ 0 & 0 & 0 & \frac{1}{\lambda^2} \end{pmatrix}, \quad \lambda \in R^\times, \mu \in R, \tag{2}$$

and we can check that any f whose matrix is $w(\lambda, \mu)$ is an automorphism of B . Also the formula

$$w(\lambda, \mu)w(\lambda', \mu') = w(\lambda\lambda', \mu' + \mu/\lambda')$$

gives that

$$\text{aut}(B) = \{w(\lambda, \mu) : \lambda \in R^\times, \mu \in R\} \cong \begin{pmatrix} 1 & R \\ 0 & R^\times \end{pmatrix}$$

being the isomorphism $w(\lambda, \mu) \mapsto \begin{pmatrix} 1 & \mu \\ 0 & \lambda^{-1} \end{pmatrix}$. So $\text{aut}(B)$ is isomorphic to the group $\text{Aff}_2(R)$ of all invertible affine transformations of the

affine plane $\mathbb{A}_2(R)$. In [22] it is proved this result in the particular case that R is a field \mathbb{F} of characteristic zero.

- (B) $\frac{1}{3} \in R$ but $2R = 0$. We have $f_1^1 = 1, f_1^2 = 0$ and $(f_1^3)^2 = 0$. Recall also that $f_4^i = 0$ for $i \neq 4$. Taking into account the multiplication table, which is the same that the multiplication of S_2 when $\text{char}(\mathbb{F}) = 2$, we deduce (from $e_3e_4 = 0$) that $f_3^1e_4 + f_3^2e_3 = 0$ so $f_3^1 = f_3^2 = 0$. Consequently we have the same pattern (1) for the matrix of the automorphism. So f_2^2, f_3^3, f_4^4 are invertible. Then we deduce $f_4^4 = (f_3^3)^2$ as in the previous case. Also following the argument in the previous case we get

$$f_2^2f_3^3 = 1, f_1^3 = 0, f_1^4 = f_3^3f_2^3.$$

Now, from $e_2^2 = 0$ we get:

$$(f_2^1)^2 + f_2^2f_2^3 = 0, f_2^2f_2^4 - f_2^3f_2^1 = 0, (f_2^3)^2 + f_2^1f_2^4 = 0.$$

If we put $\lambda = f_2^2, \mu = f_2^1$ we have $f_2^3 = (f_2^1)^2/f_2^2 = \mu^2/\lambda$ implying $f_1^4 = \mu^2/\lambda^2$. Then $f_3^3 = 1/\lambda$. On the other hand, since $e_2e_4 = e_3$ we have $f_2^ie_4f_4^4e_4 = f_3^3e_3 + f_3^4e_4$ which gives

$$f_2^1f_4^4e_4 + f_2^2f_4^4e_3 = f_3^3e_3 + f_3^4e_4, \text{ that is, } f_2^1f_4^4 = f_3^4 \text{ and } f_2^2f_4^4 = f_3^3.$$

So $f_3^4 = \mu/\lambda^2$ and $f_4^4 = 1/\lambda^2$. The matrix of f is

$$w_2(\lambda, \mu) = \begin{pmatrix} 1 & 0 & 0 & \frac{\mu^2}{\lambda^2} \\ \mu & \lambda & \frac{\mu^2}{\lambda} & \frac{\mu^3}{\lambda^2} \\ 0 & 0 & \frac{1}{\lambda} & \frac{\mu}{\lambda^2} \\ 0 & 0 & 0 & \frac{1}{\lambda^2} \end{pmatrix}, \quad \lambda \in R^\times, \mu \in R, \quad (3)$$

and reciprocally: if the matrix of f relative to the basis $\{e_i\}$ is as above, then f is an automorphism of B . We also have the formula $w_2(\lambda, \mu)w_2(\lambda', \mu') = w_2(\lambda\lambda', \mu + \lambda\mu')$ which gives an isomorphism

$$\text{aut}(B) = \{w_2(\lambda, \mu) : \lambda \in R^\times, \mu \in R\} \cong \begin{pmatrix} 1 & R \\ 0 & R^\times \end{pmatrix}$$

being the isomorphism the given by $w_2(\lambda, \mu) \mapsto \begin{pmatrix} 1 & \mu \\ 0 & \lambda \end{pmatrix}$. So again $\text{aut}(B)$ is isomorphic to the group $\text{Aff}_2(R)$ of all invertible affine transformations of the affine plane $\mathbb{A}_2(R)$.

- (C) $3R = 0$. As in previous cases $f(e_4) = f_4^4e_4$ so $f_4^i = 0$ except for $i = 4$. Imposing the condition $f(e_1^2) = f(e_1)^2$ we get $f_1^2 = f_1^4 = 0$. Imposing $f(e_1e_3) = f(e_1)f(e_3)$ we get $f_3^2 = f_3^4 = 0$ and from $f(e_2e_4) = f(e_2)f(e_4)$ we get $f_3^1 = 0$ and $f_3^3 = f_2^2f_4^4$. Taking into account these values, from $f(e_3e_1) = f(e_3)f(e_1)$ we get $f_1^1 = 1$. Aplying now $f(e_3e_2) = f(e_3)f(e_2)$ gives $f_1^3 = -f_2^1f_3^3$ and $f_4^4 = 1/(f_2^2)^2$. Using again $f(e_1e_2) = f(e_1)f(e_2)$ we get $f_2^3 = (f_2^1)^2/f_2^2$. Finally, equation $f(e_2)^2 = 0$ gives $f_2^4 = f_2^1f_3^3/f_2^2 = (f_2^1)^3/(f_2^2)^2$. If

we do $\lambda = f_2^2, \mu = f_2^1$ we have the matrix

$$w_3(\lambda, \mu) = \begin{pmatrix} 1 & 0 & -\frac{\mu}{\lambda} & 0 \\ \mu & \lambda & \frac{\mu^2}{\lambda} & \frac{\mu^3}{\lambda^2} \\ 0 & 0 & \frac{1}{\lambda} & 0 \\ 0 & 0 & 0 & \frac{1}{\lambda^2} \end{pmatrix} \tag{4}$$

so that the equality $w_3(\lambda, \mu)w_3(\lambda', \mu') = w_3(\lambda\lambda', \mu + \lambda\mu')$ holds. Then $\text{aut}(B) = \{w_3(\lambda, \mu) : \lambda \in R^\times, \mu \in R\} \cong \begin{pmatrix} 1 & R \\ 0 & R^\times \end{pmatrix}$, the isomorphism being $w_3(\lambda, \mu) \mapsto \begin{pmatrix} 1 & \mu \\ 0 & \lambda \end{pmatrix}$. So again $\text{aut}(B)$ is isomorphic to the group $\text{Aff}_2(R)$ of all invertible affine transformations of the affine plane $\mathbb{A}_2(R)$.

Thus we claim:

Proposition 8. *Let R be a commutative associative unital ring and B the free R -module with basis $\{e_i\}_{i=1}^4$ endowed with an R -algebra structure whose multiplication algebra is that of table of multiplication of S_2 . Then if $\frac{1}{2}, \frac{1}{3} \in R$; or $\frac{1}{3} \in R, 2R = 0$; or $3R = 0$, we have $\text{aut}(B) \cong \text{Aff}_2(R)$ the affine group of $\mathbb{A}_2(R)$. The precise description of $\text{aut}(B)$ is given in formulae (2), (3), (4).*

Now fix an arbitrary field \mathbb{F} and let S_2 be the \mathbb{F} -algebra introduced in the table of multiplication of S_2 . We can describe the affine group scheme $\mathbf{aut}(S_2)$. Denote by $\text{Alg}_{\mathbb{F}}$ the category of associative, commutative and unital \mathbb{F} -algebras and by Grp the category of groups. Then $\mathbf{aut}(S_2)$ is the group functor $\mathbf{aut}(S_2) : \text{Alg}_{\mathbb{F}} \rightarrow \text{Grp}$ such that $R \mapsto \text{aut}((S_2)_R)$ (where $(S_2)_R := S_2 \otimes_{\mathbb{F}} R$ is the scalar extension algebra). If $\text{ch}(\mathbb{F}) \neq 2, 3$ then $\frac{1}{2}, \frac{1}{3} \in R$ for any $R \in \text{Alg}_{\mathbb{F}}$. If $\text{ch}(\mathbb{F}) = 2$ then $2R = 0$ for any $R \in \text{Alg}_{\mathbb{F}}$ but $\frac{1}{3} \in R$. Finally, if $\text{ch}(\mathbb{F}) = 3$ then $3R = 0$. So in any case we can apply Proposition 8 to $R \in \text{Alg}_{\mathbb{F}}$ to compute the affine group scheme $\mathbf{aut}(S_2)$. Denote by $\mathbf{Aff}_2 : \text{Alg}_{\mathbb{F}} \rightarrow \text{Grp}$ the group functor such that $R \mapsto \text{Aff}_2(R) = \begin{pmatrix} 1 & R \\ 0 & R^\times \end{pmatrix}$. Then we claim

Theorem 9. *For an arbitrary field \mathbb{F} , there is an isomorphism of group schemes*

$$\mathbf{aut}(S_2) \cong \mathbf{Aff}_2.$$

We recall that the isomorphism condition between group functors is that there is a collection of group isomorphisms $\tau_R : \text{aut}((S_2)_R) \cong \text{Aff}_2(R)$ such that when $\alpha : R \rightarrow S$ is an \mathbb{F} -algebra homomorphism, the following squares commute:

$$\begin{array}{ccc} \text{aut}((S_2)_R) & \xrightarrow{\tau_R} & \text{Aff}_2(R) \\ \alpha_1 \downarrow & & \downarrow \alpha_2 \\ \text{aut}((S_2)_S) & \xrightarrow{\tau_S} & \text{Aff}_2(S) \end{array}$$

where each α_i ($i = 1, 2$) is given by applying the corresponding functor $\mathbf{aut}(S_2)$ or \mathbf{Aff}_2 to the homomorphism α . Consider now the \mathbb{F} -algebra of

dual numbers $\mathbb{F}(\epsilon)$. Recall that for an algebraic group $\mathcal{G} \subset \mathbf{GL}(V)$ (with V a finite-dimensional \mathbb{F} -vector space), its Lie algebra is $\mathbf{lie}(\mathcal{G}) = \{d \in \mathbf{gl}(V) : 1 + \epsilon d \in \mathcal{G}(\mathbb{F}(\epsilon))\}$. Thus $\mathbf{lie}(\mathbf{aut} S_2) \cong \mathbf{lie}(\mathbf{Aff}_2) \cong \mathbf{aff}_2(\mathbb{F})$ where

$$\mathbf{aff}_2(\mathbb{F}) = \left\{ \begin{pmatrix} 0 & \mu \\ 0 & \lambda \end{pmatrix} : \lambda, \mu \in \mathbb{F} \right\}.$$

As a corollary of Proposition 9 we have


Corollary 10. *For an arbitrary field we have:*

$$\mathfrak{Der}(S_2) \cong \mathbf{aff}_2(\mathbb{F}).$$


2.2. Conservative algebra W_2 . Consider now the six-dimensional \mathbb{F} -algebra W_2 whose multiplication algebra is given in the following table

	e_1	e_2	e_3	e_4	e_5	e_6
e_1	$-e_1$	$-3e_2$	e_3	$3e_4$	$-e_5$	e_6
e_2	$3e_2$	0	$2e_1$	e_3	0	$-e_5$
e_3	$-2e_3$	$-e_1$	$-3e_4$	0	e_6	0
e_4	0	0	0	0	0	0
e_5	$-2e_1$	$-3e_2$	$-e_3$	0	$-2e_5$	$-e_6$
e_6	$2e_3$	e_1	$3e_4$	0	$-e_6$	0

If \mathbb{F} is of characteristic $\neq 2, 3$, the graph of B relative to the basis of the e_i 's is



which is, of course, strongly connected. In the case $\text{char}(\mathbb{F}) = 3$ the graph is not strongly connected:



There is an ideal $I := \bigoplus_{i \neq 2} \mathbb{F}e_i$ so that $W_2 = I \oplus \mathbb{F}e_2$ and again $W_2/I \cong \mathbb{F}e_2$ is a zero-product algebra.

Remark 11. Recall that the Jacobson radical $\text{rad}(A)$ of a unital algebra A contains every nilpotent ideal of A . On the other hand if $I \triangleleft A$ and A/I is semisimple, then $\text{rad}(A) \subset I$.

Theorem 12. *For a ground field \mathbb{F} of characteristic not 3, we have $\mathcal{M}(W_2) = \text{End}_{\mathbb{F}}(W_2)$, hence W_2 is simple. In the characteristic 3 case, W_2 has a five dimensional ideal $I = \oplus_{i \neq 2} \mathbb{F}e_i$ and $W_2/I \cong \mathbb{F}e_2$ has zero product. The multiplication algebra $\mathcal{M} = \mathcal{M}(W_2)$ is 20-dimensional, its radical $\text{rad}(\mathcal{M})$ is 12-dimensional and $\mathcal{M}/\text{rad}(\mathcal{M}) = M_2(\mathbb{F}) \oplus M_2(\mathbb{F})$.*

Proof. We will apply Proposition 5 repeatedly.

- (A) Assume first that $\text{char}(\mathbb{F}) \neq 2, 3$. Then $L_{e_2}^3 = 6E_{42}$ hence $E_{4k} \in \mathcal{M} := \mathcal{M}(W_2)$ for any k (take into account Remark 6). Since $L_{e_3}^3 = -6E_{24}$ we have $E_{2k} \in \mathcal{M}$ for any k . Also $L_{e_2}^2 L_{e_5}^3 = -6E_{32}$ hence $E_{3k} \in \mathcal{M}$ for every k . Since $R_{e_2}^2 = 3E_{32} - 3E_{62}$ we conclude $E_{6k} \in \mathcal{M}$ for every k . Furthermore $L_{e_2} = 3E_{12} + 2E_{31} + E_{43} - E_{65}$ hence $E_{1k} \in \mathcal{M}$ for any k . And finally $R_{e_2} = -3E_{12} - E_{31} - 3E_{52} + E_{61}$ which implies $E_{5k} \in \mathcal{M}$ for every k . Thus $\mathcal{M} = \text{End}_{\mathbb{F}}(W_2)$.
- (B) When the ground field has characteristic 2 we take into account:

$$\begin{array}{llll}
 R_{e_2}^2 L_{e_2} & = & E_{42} & \text{implying } E_{4k} \in \mathcal{M} \text{ for any } k. \\
 R_{e_2} L_{e_2} & = & E_{41} + E_{62} & \text{implying } E_{6k} \in \mathcal{M} \text{ for any } k. \\
 L_{e_2} & = & E_{12} + E_{43} + E_{65} & \text{implying } E_{1k} \in \mathcal{M} \text{ for any } k. \\
 R_{e_4} & = & E_{14} + E_{23} & \text{implying } E_{2k} \in \mathcal{M} \text{ for any } k. \\
 L_{e_5} & = & E_{22} + E_{33} + E_{66} & \text{implying } E_{3k} \in \mathcal{M} \text{ for any } k. \\
 L_{e_1} & = & \sum_1^6 E_{ii} & \text{implying } E_{5k} \in \mathcal{M} \text{ for any } k.
 \end{array}$$

- (C) In case $\text{char}(\mathbb{F}) = 3$ we have

$$\begin{array}{ll}
 L_{e_1} = -E_{11} + E_{33} - E_{55} + E_{66}, & R_{e_1} = -E_{11} + E_{33} + E_{51} - E_{63}, \\
 L_{e_2} = -E_{31} + E_{43} - E_{65}, & R_{e_2} = E_{61} - E_{31}, \\
 L_{e_3} = E_{13} - E_{21} + E_{56}, & R_{e_3} = E_{13} - E_{21} - E_{53}, \\
 L_{e_4} = 0, & R_{e_4} = E_{23}, \\
 L_{e_5} = E_{11} - E_{33} + E_{55} - E_{66}, & R_{e_5} = -E_{15} + E_{36} + E_{55} - E_{66}, \\
 L_{e_6} = -E_{13} + E_{21} - E_{56}, & R_{e_6} = E_{16} - E_{25} - E_{56}.
 \end{array}$$

A basis for \mathcal{M} is given by the set of matrices:

$$\begin{array}{cccc}
 E_{11} + E_{55}, & E_{33} + E_{66}, & E_{31} + E_{65}, & E_{13} + E_{56}, \\
 E_{11} - E_{51}, & E_{33} - E_{63}, & E_{31} - E_{61}, & E_{13} - E_{53}, \\
 E_{15} - E_{55}, & E_{36} - E_{66}, & E_{16} - E_{56}, & E_{35} - E_{65}, \\
 E_{21}, & E_{23}, & E_{25}, & E_{26}, \\
 E_{41}, & E_{43}, & E_{45}, & E_{46}.
 \end{array}$$

We have computed again the radical \mathcal{M}^\perp of its trace form $\langle f, g \rangle := \text{Tr}(fg)$ and it is 12-dimensional ideal. More precisely

$$\begin{aligned}
 \mathcal{M}^\perp & := \text{rad}(\langle \cdot, \cdot \rangle) = \\
 & \mathbb{F}E_{21} \oplus \mathbb{F}E_{23} \oplus \mathbb{F}E_{25} \oplus \mathbb{F}E_{26} \oplus \mathbb{F}E_{41} \oplus \mathbb{F}E_{43} \oplus \mathbb{F}E_{45} \oplus \mathbb{F}E_{46} \oplus \\
 & \mathbb{F}(E_{11} + E_{15} - E_{51} + 2E_{55}) \oplus \mathbb{F}(E_{13} + E_{16} - E_{53} + 2E_{56}) \oplus \\
 & \mathbb{F}(E_{31} + E_{35} - E_{61} + 2E_{65}) \oplus \mathbb{F}(E_{33} + E_{36} - E_{63} + 2E_{66})
 \end{aligned}$$

and $(\mathcal{M}^\perp)^2 = \mathbb{F}(E_{43} + E_{46}) \oplus \mathbb{F}(E_{41} + E_{45}) + \mathbb{F}(E_{23} + E_{26}) + \mathbb{F}(E_{21} + E_{25})$ being $(\mathcal{M}^\perp)^4 = 0$. Since \mathcal{M}^\perp is nilpotent, $\text{rad}(\mathcal{M}) \supset \mathcal{M}^\perp$ (see Remark 11). Define next the subspace S of \mathcal{M} whose basis is $\{e_{ij}\}_{i,j=1}^2 \sqcup \{u_{ij}\}_{i,j=1}^2$ given by

$$\begin{aligned} e_{1,1} &= E_{11} + E_{15}, & e_{2,2} &= E_{33} + E_{36}, \\ e_{1,2} &= E_{13} + E_{16}, & e_{2,1} &= E_{31} + E_{35}, \\ u_{1,1} &= -E_{15} + E_{55}, & u_{2,2} &= -E_{36} + E_{66}, \\ u_{1,2} &= E_{16} - E_{56}, & u_{2,1} &= E_{35} - E_{65}. \end{aligned}$$

If δ_{ij} denotes the Kronecker delta, it is easy to check that

$$e_{ij}e_{kl} = \delta_{jk}e_{il}, \quad u_{ij}u_{kl} = \delta_{jk}u_{il} \quad \text{and} \quad e_{ij}u_{kl} = u_{kl}e_{ij} = 0$$

for any $i, j, k, l \in \{1, 2\}$. Thus $S \cong M_2(\mathbb{F}) \oplus M_2(\mathbb{F})$ and furthermore $\mathcal{M} = \mathcal{M}^\perp \oplus S$. Thus $\mathcal{M}/\mathcal{M}^\perp \cong M_2(\mathbb{F}) \oplus M_2(\mathbb{F})$ is semisimple which implies $\text{rad}(\mathcal{M}) \subset \mathcal{M}^\perp$. So $\text{rad}(\mathcal{M}) = \mathcal{M}^\perp$. □

2.2.1. Automorphisms of W_2 . In this section we work over a commutative ring R and denote $W_2(R)$ the R -algebra $\bigoplus_{i=1}^6 Re_i$ where the multiplication table of the e_i 's is that of the multiplication table of W_2 given above. If we take a generic element $w = \sum_{i=1}^6 \lambda_i e_i \in W_2(R)$ and compute the matrix of L_w relative to the basis of the e_i 's we obtain:

$$\begin{pmatrix} -\lambda_1 - 2\lambda_5 & 3\lambda_2 & 2\lambda_6 - 2\lambda_3 & 0 & 0 & 0 \\ \lambda_6 - \lambda_3 & -3\lambda_1 - 3\lambda_5 & 0 & 0 & 0 & 0 \\ 2\lambda_2 & 0 & \lambda_1 - \lambda_5 & 3\lambda_6 - 3\lambda_3 & 0 & 0 \\ 0 & 0 & \lambda_2 & 3\lambda_1 & 0 & 0 \\ 0 & 0 & 0 & 0 & -\lambda_1 - 2\lambda_5 & \lambda_3 - \lambda_6 \\ 0 & 0 & 0 & 0 & -\lambda_2 & \lambda_1 - \lambda_5 \end{pmatrix}.$$

If f is any element in $\text{aut}(W_2(R))$ we know $L_{f(w)} = fL_w f^{-1}$ hence the characteristic polynomial of L_w is invariant under automorphism of $W_2(R)$. So the coefficients of that polynomial are invariants and we list here:

$$\begin{aligned} \ell_1(w) &:= 9\lambda_5, \\ \ell_2(w) &:= -11\lambda_1^2 - 11\lambda_5\lambda_1 + 31\lambda_5^2 + 11\lambda_2\lambda_3 - 11\lambda_2\lambda_6, \\ \ell_3(w) &:= -3\lambda_5 (22\lambda_1^2 + 22\lambda_5\lambda_1 - 17\lambda_5^2 - 22\lambda_2\lambda_3 + 22\lambda_2\lambda_6), \\ \ell_4(w) &:= 19\lambda_1^4 + 38\lambda_5\lambda_1^3 - 120\lambda_5^2\lambda_1^2 - 38\lambda_2\lambda_3\lambda_1^2 + 38\lambda_2\lambda_6\lambda_1^2 - 139\lambda_5^3\lambda_1 - \\ &\quad 38\lambda_2\lambda_3\lambda_5\lambda_1 + 38\lambda_2\lambda_5\lambda_6\lambda_1 + 40\lambda_5^4 + \\ &\quad 19\lambda_2^2\lambda_3^2 + 139\lambda_2\lambda_3\lambda_5^2 + 19\lambda_2^2\lambda_6^2 - 139\lambda_2\lambda_5^2\lambda_6 - 38\lambda_2^2\lambda_3\lambda_6, \\ \ell_5(w) &:= 3\lambda_5 (\lambda_1^2 + \lambda_5\lambda_1 - 2\lambda_5^2 - \lambda_2\lambda_3 + \lambda_2\lambda_6) \cdot \\ &\quad (19\lambda_1^2 + 19\lambda_5\lambda_1 - 2\lambda_5^2 - 19\lambda_2\lambda_3 + 19\lambda_2\lambda_6), \\ \ell_6(w) &:= -9 (\lambda_1^2 + \lambda_5\lambda_1 - \lambda_2\lambda_3 + \lambda_2\lambda_6) \cdot \\ &\quad (\lambda_1^2 + \lambda_5\lambda_1 - 2\lambda_5^2 - \lambda_2\lambda_3 + \lambda_2\lambda_6)^2. \end{aligned}$$

All these polynomial remain invariant under automorphism but we are using only the first one: ℓ_1 .

Lemma 13. *If an element $x \in W_2(R) \setminus \{0\}$ with $\ell_1(x) = 9$ satisfies $x^2 = -2x$ and $\text{Tor}_3(R) = 0$ then $x = se_4 + e_5 + re_6$ for some $r, s \in R$ such that $2s = 0$. In particular if $\frac{1}{2} \in R$ we have $x = e_5 + re_6$.*

Proof. Take $x = x_i e_i$ where $x_i \in R$. Since $\ell_1(x) = 9$ we have $x_5 = 1$. By using the multiplication table, from the equality $x^2 = -2x$ we also obtain

$$\begin{aligned} -x_1^2 + x_2x_3 + x_2x_6 &= 0, \\ x_2 &= 0, \\ -x_1x_3 + x_3 + x_2x_4 + 2x_1x_6 &= 0, \\ -3x_3^2 + 3x_6x_3 + 3x_1x_4 + 2x_4 &= 0, \\ -x_1 - x_2x_6 &= 0, \\ x_3 + x_1x_6 &= 0, \end{aligned}$$

which can be summarized in $x_1 = x_2 = x_3 = 2x_4 = 0$. Thus taking $s = x_4$ and $r = x_6$ the Lemma is proved. \square

If $\theta \in \text{aut}(W_2(R))$ then $\theta(e_5)^2 = -2\theta(e_5)$ and $\ell_1(\theta(e_5)) = \ell_1(e_5) = 9$. Hence Lemma 13 implies that if $\text{Tor}_3(R) = 0$ then $\theta(e_5) = se_4 + e_5 + re_6$ with $2s = 0$. In case $\text{Tor}_2(R) = 0$ we have $\theta(e_5) = e_5 + re_6$.

Lemma 14. *Let again $\text{Tor}_3(R) = 0$. If $x, y \in W_2(R)$ are linearly independent with $\ell_1(x) = 9, \ell_1(y) = 0$, and they satisfies $x^2 = -2x, xy = yx = -y, y^2 = 0$ then in case $\frac{1}{2} \in R$ we have $x = e_5 + re_6$ and $y = te_6$ for some $r, t \in R$. If $\text{Tor}_2(R) = R$ we can only conclude that $x = se_4 + e_5 + re_6$ and $y = te_6$ for some $t \in R$.*

Proof. From Lemma 13 we know that if $\frac{1}{2} \in R$ we have $x = e_5 + re_6$. Now writing $y = \sum \mu_i e_i$ (with $\mu_5 = 0$) and imposing $xy = yx = -y$ we get the equations:

$$\begin{aligned} 2r\mu_1 &= 0, & -3\mu_2 &= 0, \\ -2\mu_2 &= 0, & r\mu_2 - 2\mu_1 &= 0, \\ r\mu_2 - \mu_1 &= 0, & \mu_1 + r\mu_2 &= 0, \\ -r\mu_1 - \mu_3 &= 0, & 2r\mu_1 - \mu_3 &= 0, \\ 3r\mu_3 &= 0, & 3r\mu_3 + \mu_4 &= 0. \end{aligned}$$

If $\frac{1}{2} \in R$ or $\text{Tor}_2(R) = R$, the first row of equations above implies $\mu_2 = 0, \mu_1 = 0$. Now the second row gives $\mu_3 = \mu_4 = 0$. Therefore $y = \mu_6 e_6$. If $\text{Tor}_2(R) = R$ imposing the conditions we only get $x = se_4 + e_5 + re_6$ and $y = te_6$. \square

For any $\theta \in \text{aut}(W_2(R))$ we can apply Lemma 14 taking $x = \theta(e_5)$ and $y = \theta(e_6)$. Thus, if $\frac{1}{2} \in R$ or $\text{Tor}_2(R) = R$ we have $\theta(e_5) = e_5 + re_6$ and $\theta(e_6) = te_6$ with $r, t \in R$ and t invertible. In case $\text{Tor}_2(R) = R$ we can argue as follows: $\theta(e_2)\theta(e_6) = \theta(e_5)$ so $t\theta(e_2)e_6 = se_4 + e_5 + re_6$ with t invertible. But the image of Re_6 is the R -submodule $Re_4 \oplus Re_6$ hence $s = 0$. Consequently

Lemma 15. *Assume that $\text{Tor}_3(R) = 0$. Then if $\frac{1}{2} \in R$ or $\text{Tor}_2(R) = R$ we have $\theta(e_5) = e_5 + re_6$ and $\theta(e_6) = te_6$ with $r, t \in R$ and t invertible.*

It can be checked that the left annihilator of $W_2(R)$ is the R -submodule of all elements $a_1(e_1 + e_5) + a_3(e_3 + e_6) + a_4e_4$ such that $a_i \in R$ with $3a_1 = 0$:

$$\text{Lann}(W_2(R)) = \text{Tor}_3(R)(e_1 + e_5) \oplus R(e_3 + e_6) \oplus Re_4 \tag{5}$$

Lemma 16. *If a is in the left annihilator of $W_2(R)$ and satisfies $e_6a = e_5a = 0$ then $a \in Re_4$*

Proof. Write $a = \alpha(e_1 + e_5) + \beta(e_3 + e_6) + \gamma e_4$ with $\alpha, \beta, \gamma \in R$, $3\alpha = 0$. Then

$$0 = e_6a = 2\alpha e_3 - \alpha e_6 + 3\beta e_4 \text{ so } \alpha = 0.$$

But $0 = e_5a = -\beta e_3 - \beta e_6 + 3\beta e_4t$ and consequently also $\beta = 0$. Thus $a = \gamma e_4$. \square

In case $\text{Tor}_3(R) = 0$ and $\frac{1}{2} \in R$ or $\text{Tor}_2(R) = R$ we have proved that $\theta(e_5) = e_5 + re_6$, $\theta(e_6) = te_6$, $r, t \in R$, $t \in R^\times$. We can apply Lemma 16 taking $a = \theta(e_4)$ for any $\theta \in \text{aut}(W_2(R))$. This implies that $\theta(e_4) = se_4$ for some invertible $s \in R$. So far, when $\text{Tor}_3(R) = 0$ and either $\frac{1}{2} \in R$ or $\text{Tor}_2(R) = R$, the matrix of an automorphism of $W_2(R)$ is of the form:

$$\left(\begin{array}{ccc|cc} * & & & * & * \\ 0 & 0 & 0 & s & 0 & 0 \\ \hline & & \mathbf{0} & & 1 & r \\ & & & & 0 & t \end{array} \right), \tag{6}$$

with $r, s, t \in R$, $s, t \in R^\times$.

We now investigate the image of e_3 under automorphisms of $W_2(R)$.

Lemma 17. *Assume $x \in W_2(R)$ satisfies $xe_4 = xe_6 = 0$, $xe_5 = te_6$ ($t \in R^\times$) and $e_6x \in Re_4$. Then if $\text{Tor}_3(R) = 0$ we have $x \in Re_3 + Re_4 + Re_6$.*

Proof. If $x = \sum_{i=1}^6 x_i e_i$ then from $xe_4 = 0$ we get $x_2 = 3x_1 = 0$. But $xe_6 = 0$ gives $x_2 = 0$ as before and $x_5 = x_1$. On the other hand $xe_5 = te_6$ gives $-x_1 - 2x_5 = 0$, $x_3 - x_6 - t = 0$. So $3x_5 = 0$ and $x_3 - x_6 \in R^\times$. Also $e_6x = 2x_1e_3 - x_1e_6 + 3x_3e_4$ giving $x_1 = 0$. Thus $x = x_3e_3 + x_4e_4 + x_6e_6$. \square

Lemma 18. *Assume $\text{Tor}_3(R) = 0$ and let $u = \sum_{i=1}^6 \lambda_i e_i \in W_2(R)$ be such that $u^2 = 0$, $\ell_1(u) = 0$, $u(e_5 + re_6) = 0$, $tue_6 = -(e_5 + re_6)$ then*

- (1) $\lambda_2 \in R^\times$ and $u = \frac{\lambda_1^3}{\lambda_2^3} e_4 + \frac{\lambda_1^2}{\lambda_2^2} e_3 + \lambda_1 e_1 + \lambda_2 e_2$.
- (2) $r = -\lambda_1 t$.
- (3) $\ell_1(u(x_3e_3 + x_4e_4 + x_6e_6)) = -9\lambda_2 x_6$.

Proof. Since $0 = \ell_1(u) = 3\lambda_5$ we have $\lambda_5 = 0$. The fact that $u(e_5 + e_6) = 0$ gives $\lambda_1 + r\lambda_2 = 0 = r\lambda_1 + \lambda_3 - \lambda_6$. Since $tue_6 = -(e_5 + re_6)$ we get the equalities $t\lambda_2 = 1$ and $t\lambda_1 = -r$. Thus λ_2 is invertible. Also since $u^2 = 0$ we have $\lambda_2\lambda_6 = 0$ which implies $\lambda_6 = 0$. Then $u = \sum_{i=1}^4 \lambda_i e_i$ and

$$\ell_1(u(x_3e_3 + x_4e_4 + x_6e_6)) = -9\lambda_2x_6$$

is readily checked. Now imposing the condition that $u^2 = 0$ gives $\lambda_3 = \lambda_1^2/\lambda_2$ and $\lambda_4 = \lambda_1^3/\lambda_2^2$. \square

Assume again $\text{Tor}_3(R) = 0$, $\frac{1}{2} \in R$ and $\theta \in \text{aut}_R(W_2(R))$. We know that $\theta(e_6) = te_6$, $\theta(e_5) = e_5 + re_6$, $\theta(e_4) = se_4$. Applying Lemma 17 we know that $\theta(e_3) = x_3e_3 + x_4e_4 + x_6e_6$ for some $x_i \in R$. Applying Lemma 18 we have $\theta(e_2) = \frac{\lambda_1^3}{\lambda_2^2}e_4 + \frac{\lambda_1^2}{\lambda_2}e_3 + \lambda_1e_1 + \lambda_2e_2$ for suitable $\lambda_i \in R$ with λ_2 invertible. Furthermore, since $e_2e_3 = 2e_1$ we have $\theta(e_2)\theta(e_3) = 2\theta(e_1)$ hence

$$-9\lambda_2x_6 = \ell_1(\theta(e_2)\theta(e_3)) = 2\ell_1(\theta(e_1)) = 2\ell_1(e_1) = 0.$$

Since λ_2 is invertible we get $x_6 = 0$ (so $\theta(e_3) \in Re_3 + Re_4$). Finally

$$\theta(e_1) = -\theta(e_3)\theta(e_2) = \frac{3\lambda_1^2x_3}{\lambda_2}e_4 + 2\lambda_1x_3e_3 + \lambda_2x_3e_1.$$

In conclusion we have

Lemma 19. *If $\text{Tor}_3(R) = 0$ and either $\frac{1}{2} \in R$ or $\text{Tor}_2(R) = R$ the matrix of $\theta \in \text{aut}_R(W_2(R))$ in the basis of the e_i 's is of the form:*

$$\begin{pmatrix} \lambda_2x_3 & 0 & 2\lambda_1x_3 & \frac{3\lambda_1^2x_3}{\lambda_2} & 0 & 0 \\ \lambda_1 & \lambda_2 & \frac{\lambda_1^2}{\lambda_2} & \frac{\lambda_1^3}{\lambda_2^2} & 0 & 0 \\ 0 & 0 & x_3 & x_4 & 0 & 0 \\ 0 & 0 & 0 & s & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & -\lambda_1t \\ 0 & 0 & 0 & 0 & 0 & t \end{pmatrix} \tag{7}$$

The algebra $W_2(R)$ contains $\bigoplus_{i=1}^4 Re_i = (S_2)_R$ as a subalgebra. Also $W_2(R) \supset Re_5 \oplus Re_6$ also as a subalgebra. Then, under the assumptions of Lemma 19 we know that any automorphism $\theta \in \text{aut}_R(W_2(R))$ preserves both subalgebras $\theta((S_2)_R) = (S_2)_R$ and $\theta(Re_5 + Re_6) = Re_5 + Re_6$. The map

$$\text{aut}_R(W_2(R)) \rightarrow \text{aut}((S_2)_R)$$

such that $\theta \mapsto \theta|_{(S_2)_R}$ is a group homomorphism. In fact it is a monomorphism because in case $\theta|_{(S_2)_R} = 1$ we have $\lambda_1 = 0$ (see equation (7)). Thus if θ fixes e_1, \dots, e_4 then also $\theta(e_5) = e_5$. Moreover since $e_3e_5 = e_6$ then θ fixes also e_6 whence $\theta = 1$.

Proposition 20. *If $\frac{1}{3}, \frac{1}{2} \in R$ or $\frac{1}{3} \in R$, $2R = 0$, the map $\text{aut}_R(W_2(R)) \rightarrow \text{aut}((S_2)_R)$ such that $\theta \mapsto \theta|_{(S_2)_R}$ is a group isomorphism.*

Proof. It only remains to prove that the map is an epimorphism. So if $\frac{1}{3}, \frac{1}{2} \in R$, take an arbitrary $f \in \text{aut}_R((S)_R)$ whose matrix relative to the basis of the e_i 's is given in (2). Define next $\hat{f}: W_2(R) \rightarrow W_2(R)$ whose restriction to $(S_2)_R$ is f and $\hat{f}(e_5) = e_5 - \frac{t}{2}e_6$, $\hat{f}(e_6) = \frac{1}{\lambda}e_6$. It can be checked that $\hat{f} \in \text{aut}_R(W_2(R))$ and $\hat{f}|_{(S_2)_R} = f$. Now in case $\frac{1}{3} \in R$ and $2R = 0$ take an arbitrary $f \in \text{aut}_R((S)_R)$ whose matrix relative to the basis

of the e_i 's is given in (3). Then extend f to the automorphism \hat{f} of $(W_2)_R$ such that $\hat{f}(e_5) = e_5 + \frac{t}{\lambda}e_6$ and $\hat{f}(e_6) = \frac{1}{\lambda}e_6$. \square

If $\text{ch}(\mathbb{F}) \neq 3$ we can describe now the affine group scheme $\mathbf{aut}(W_2)$. As before $\text{Alg}_{\mathbb{F}}$ will be the category of associative, commutative and unital \mathbb{F} -algebras and Grp that of groups. Then $\mathbf{aut}(W_2)$ is the group functor $\mathbf{aut}(W_2): \text{Alg}_{\mathbb{F}} \rightarrow \text{Grp}$ such that $R \mapsto \text{aut}_R((W_2)_R)$ (as usual $(W_2)_R := W_2 \otimes_{\mathbb{F}} R$ is the scalar extension algebra). If $\text{ch}(\mathbb{F}) \neq 2, 3$ then $\frac{1}{2}, \frac{1}{3} \in R$ for any $R \in \text{Alg}_{\mathbb{F}}$. If $\text{ch}(\mathbb{F}) = 2$ then $2R = 0$ for any $R \in \text{Alg}_{\mathbb{F}}$ but $\frac{1}{3} \in R$. So in any case we can apply Proposition 20 to $R \in \text{Alg}_{\mathbb{F}}$ to compute the affine group scheme $\mathbf{aut}(W_2(R))$. Denote by $\mathbf{Aff}_2: \text{Alg}_{\mathbb{F}} \rightarrow \text{Grp}$ the group functor such that $R \mapsto \text{Aff}_2(R) = \begin{pmatrix} 1 & R \\ 0 & R^\times \end{pmatrix}$. Then we claim

Theorem 21. *For a field \mathbb{F} with $\text{ch}(\mathbb{F}) \neq 3$, there is an isomorphism of group schemes*

$$\mathbf{aut}(W_2) \cong \mathbf{Aff}_2.$$

Any automorphism f of $W_2(R)$ is of the form in (7) relative to the standard basis. We can refine its form a little. If we impose $f(e_i e_j) = f(e_i) f(e_j)$ for:

- (1) $i = 3, j = 5$ we get $x_3 = t$.
- (2) $i = 1, j = 3$ we get $x_3 \lambda_2 = 1$.
- (3) $i = 2, j = 6$ we get $\lambda_2 = 1/t$.
- (4) $i = 6, j = 3$ we get $s = t^2$.
- (5) $i = 5, j = 3$ we get $x_4 = 3t^2 \lambda_1$.

Thus the form of a general automorphism of $W_2(R)$ when $\text{Tor}_3(R) = 0$ on canonical basis is

$$w(x, t) := \begin{pmatrix} 1 & 0 & 2tx & 3t^2x^2 & 0 & 0 \\ x & \frac{1}{t} & tx^2 & t^2x^3 & 0 & 0 \\ 0 & 0 & t & 3t^2x & 0 & 0 \\ 0 & 0 & 0 & t^2 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & -tx \\ 0 & 0 & 0 & 0 & 0 & t \end{pmatrix} \tag{8}$$

So $\text{aut}(W_2(R)) = \{w(x, t): x \in R, t \in R^\times\}$ and we can observe that $w(0, 1)$ is the identity and that

$$w(x, t)w(x', t') = w\left(x + \frac{x'}{t}, tt'\right) \text{ and } w(x, t)^{-1} = w(-tx, t^{-1})$$

for any x, x', t, t' .

2.2.2. The case of characteristic 3. In order to investigate the case of characteristic 3 we will need to note that if $3R = 0$ we have

$$W_2(R)^2 = Re_1 \oplus Re_3 \oplus Re_5 \oplus Re_6.$$

So for $i = 1, 3, 5, 6$ and any $\theta \in \text{aut}_R(W_2(R))$ we have $\theta(e_i) \in Re_1 \oplus Re_3 \oplus Re_5 \oplus Re_6$. Now $\theta(e_5) = \lambda_1 e_1 + \lambda_3 e_3 + \lambda_5 e_5 + \lambda_6 e_6$ and since $e_5^2 = e_5$, applying θ we get $2\lambda_1^2 + \lambda_1 \lambda_5 + 2\lambda_1 = 0$. Now we will use the invariant ℓ_2

(see section 2.2.1). We have $1 = \ell_2(e_5) = \ell_2(\theta(e_5)) = \lambda_1^2 + \lambda_5\lambda_1 + \lambda_5^2$ so that $\lambda_1^2 + \lambda_5\lambda_1 + \lambda_5^2 = 1$. Now from

$$\begin{cases} 2\lambda_1^2 + \lambda_5\lambda_1 + 2\lambda_1 = 0 \\ \lambda_1^2 + \lambda_5\lambda_1 + \lambda_5^2 = 1 \end{cases}$$

we get $2\lambda_1\lambda_5 + 2\lambda_1 + \lambda_5^2 = 1$. Also from the equality $e_5^2 = e_5$ applying θ we get $\lambda_5^2 + 2\lambda_1\lambda_5 + 2\lambda_5 = 0$ so that $\lambda_5 = 1 + \lambda_1$. Thus we can write

$$\theta(e_5) = \lambda_1 e_1 + \lambda_3 e_3 + (1 + \lambda_1)e_5 + \lambda_6 e_6. \quad (9)$$

On the other hand by equation (5) we can write

$$\theta(e_4) = \alpha(e_1 + e_5) + \beta(e_3 + e_6) + \gamma e_4 \text{ and since } \theta(e_5)\theta(e_4) = 0$$

after an easy calculation we get $\alpha = \beta = 0$ so that $\theta(e_4) = t e_4$ (with $t \in R^\times$). Now we put $\theta(e_2) = \sum_{i=1}^6 y_i e_i$ and $\theta(e_3) = z_1 e_1 + z_3 e_3 + z_5 e_5 + z_6 e_6$, for scalars $y_i, z_j \in R$, and given that $\theta(e_2)\theta(e_4) = \theta(e_3)$ we get

$$t y_2 = z_3, z_1 = z_5 = z_6 = 0$$

so that $\theta(e_3) = t y_2 e_3$. Consequently $y_2 \in R^\times$. Now writing

$$\theta(e_1) = x_1 e_1 + x_3 e_3 + x_5 e_5 + x_6 e_6,$$

since $\theta(e_3)\theta(e_1) = \theta(e_3)$ we get $x_1 = 1$ and $x_5 = 0$ and so

$$\theta(e_1) = e_1 + x_3 e_3 + x_6 e_6.$$

Since $\theta(e_1)^2 = 2\theta(e_1)$ after expanding the corresponding equation we get $x_6 = 0$ so $\theta(e_1) = e_1 + x_3 e_3$. Now $\theta(e_1)\theta(e_2) = 0$ gives

$$y_1 = -x_3 y_2, y_3 = -x_3 y_1, y_5 = 0, y_6 = 0.$$

Thus $\theta(e_2) = -x_3 y_2 e_1 + y_2 e_2 + x_3^2 y_2 e_3 + y_4 e_4$. Also $\theta(e_1)\theta(e_5) = 2\theta(e_5)$ which implies

$$\lambda_1 x_3 - \lambda_3 = 0 = -\lambda_6 + \lambda_1 x_3 + x_3.$$

Moreover, $\theta(e_1)\theta(e_6) = \theta(e_6)$ and if we write $\theta(e_6) = \mu_1 e_1 + \mu_3 e_3 + \mu_5 e_5 + \mu_6 e_6$ we get

$$\mu_1 = \mu_5 = 0.$$

Also $\theta(e_2)\theta(e_3) = 2\theta(e_1)$ which implies $t = 1/y_2^2$. On the other hand the equality $\theta(e_2)^2 = 0$ gives $y_2(y_4 + x_3^3 y_2) = 0$ and the invertibility of y_2 implies $y_4 = -x_3^3 y_2$. Finally since $\theta(e_2)\theta(e_6) = 2\theta(e_5)$ we get $\lambda_1 = y_2 \mu_3$ and $\mu_6 = \frac{\mu_3 y_2 + 1}{y_2}$. Assambling all of this together we get to the matrix of and automorphism θ of $W_2(R)$:

$$\begin{pmatrix} 1 & 0 & x_3 & 0 & 0 & 0 \\ -x_3 y_2 & y_2 & x_3^2 y_2 & -x_3^3 y_2 & 0 & 0 \\ 0 & 0 & \frac{1}{y_2} & 0 & 0 & 0 \\ 0 & 0 & 0 & \frac{1}{y_2^2} & 0 & 0 \\ \mu_3 y_2 & 0 & \mu_3 x_3 y_2 & 0 & \mu_3 y_2 + 1 & \mu_3 x_3 y_2 + x_3 \\ 0 & 0 & \mu_3 & 0 & 0 & \frac{\mu_3 y_2 + 1}{y_2} \end{pmatrix}$$

whose determinant is $\frac{(\mu_3 y_2 + 1)^2}{y_2^3}$. Hence $\mu_3 y_2 + 1 \in R^\times$. So using the parameters $a = 1/y_2, c = x_3, b = \mu_3/a + 1$ the matrix of a general automorphism θ (in the basis of the e_i 's) is

$$M_{a,b,c} := \begin{pmatrix} 1 & 0 & c & 0 & 0 & 0 \\ -\frac{c}{a} & \frac{1}{a} & \frac{c^2}{a} & -\frac{c^3}{a} & 0 & 0 \\ 0 & 0 & a & 0 & 0 & 0 \\ 0 & 0 & 0 & a^2 & 0 & 0 \\ b-1 & 0 & (b-1)c & 0 & b & bc \\ 0 & 0 & a(b-1) & 0 & 0 & ab \end{pmatrix} \tag{10}$$

where $a \in R^\times, c \in R$ and $b = y_2 \mu_3 + 1$ hence $b \in R^\times$. In fact the set $\{M_{a,b,c}: a, b \in R^\times, c \in R\}$ is a group relative to matrix multiplications and obeys the rule:

$$M_{a,b,c} M_{a',b',c'} = M_{aa',bb',a'c+c'}$$

hence its identity is $M_{1,1,0}$ and also $M_{a,b,c}^{-1} = M_{a^{-1},b^{-1},-ca^{-1}}$. Thus

$$\text{aut}_R(W_2(R)) \cong \{M_{a,b,c}: a, b \in R^\times, c \in R\} \tag{11}$$

Modulo the above identification we can see that the subset $\{M_{a,1,c}: a \in R^\times, c \in R\}$ is a normal subgroup of $\text{aut}_R(W_2(R))$ and it is isomorphic to $\text{Aff}_2(R)$. Of course the quotient group is isomorphic to the multiplicative group:

$$\text{aut}(W_2(R))/\text{Aff}_2(R) \cong R^\times.$$

Let us compute the center $Z(\text{aut}_R(W_2(R)))$, in we consider the equality $M_{a,b,c} M_{x,y,z} = M_{x,y,z} M_{a,b,c}$ for a fixed triple $(a, b, c) \in R^\times \times R^\times \times R$ and an arbitrary one $(x, y, z) \in R^\times \times R^\times \times R$, we find that $cx + z = az + c$ hence taking $x = 1$ we get $z = az$ for any $z \in R$. So $a = 1$ and this implies $cx = c$ for any $x \in R^\times$. Thus $c(-x) = c$ also. Consequently $2c = 0$ and since $3R = 0$ this implies $c = 0$. Then

$$Z(\text{aut}_R(W_2(R))) = \{M_{1,b,0}: b \in R^\times\} \cong R^\times.$$

Furthermore we have a decomposition $M_{x,y,z} = M_{1,y,0} M_{x,1,z}$ for any $M_{x,y,z}$. So we have an isomorphism

$$\text{aut}_R(W_2(R)) \cong R^\times \times \text{Aff}_2(R)$$

$$M_{x,y,z} \mapsto \left(y, \begin{pmatrix} 1 & z \\ 0 & x \end{pmatrix} \right)$$

induced by the decomposition of $\text{aut}_R(W_2(R))$ as a direct product of groups. Summarizing the previous results we have:

Theorem 22. *If $3R = 0$ we have an isomorphism $\text{aut}_R(W_2(R)) \cong R^\times \times \text{Aff}_2(R)$ where $Z(\text{aut}_R(W_2(R)))$ corresponds with the first factor R^\times . Modulo the above identification, both subgroups R^\times and $\text{Aff}_2(R)$ are normal subgroups. The general form of an element $M_{a,b,c} \in \text{aut}_R(W_2(R))$ is in equation (10) relative to the basis of the e_i 's.*

2.3. Conservative algebra $W(2)$. A multiplication on the 2-dimensional vector space V_2 is defined by a $2 \times 2 \times 2$ matrix. Their classification was given in many papers (see, for example, [23, 25]). Let us consider the space $W(2)$ of all multiplications on the 2-dimensional space V_2 with a basis v_1, v_2 . The definition of the multiplication \cdot on the algebra $W(2)$ can be found in Introduction (see also [15, 20, 22]). Namely, we fix the vector $v_1 \in V_2$ and define

$$(A \cdot B)(x, y) = A(v_1, B(x, y)) - B(A(v_1, x), y) - B(x, A(v_1, y))$$

for $x, y \in V_2$ and $A, B \in W(2)$. The algebra $W(2)$ is conservative [15].

Let us consider the multiplications α_{ij}^k ($i, j, k = 1, 2$) on V_2 defined by the formula $\alpha_{ij}^k(v_t, v_l) = \delta_{it}\delta_{jl}v_k$ for all t, l . It is easy to see that $\{\alpha_{ij}^k | i, j, k = 1, 2\}$ is a basis of the algebra $W(2)$. The multiplication table of $W(2)$ in this basis is given in [20]. In this work we use another basis for the algebra $W(2)$. Let introduce the notation


$$\begin{aligned} e_1 &= \alpha_{11}^1 - \alpha_{12}^2 - \alpha_{21}^2, & e_2 &= \alpha_{11}^2, \\ e_3 &= \alpha_{22}^2 - \alpha_{12}^1 - \alpha_{21}^1, & e_4 &= \alpha_{22}^1, \\ e_5 &= 2\alpha_{11}^1 + \alpha_{12}^2 + \alpha_{21}^2, & e_6 &= 2\alpha_{22}^2 + \alpha_{12}^1 + \alpha_{21}^1, \\ e_7 &= \alpha_{12}^1 - \alpha_{21}^1, & e_8 &= \alpha_{12}^2 - \alpha_{21}^2. \end{aligned}$$

It is easy to see that the multiplication table of $W(2)$ in the basis e_1, \dots, e_8 is the one in following figure:

	e_1	e_2	e_3	e_4	e_5	e_6	e_7	e_8
e_1	$-e_1$	$-3e_2$	e_3	$3e_4$	$-e_5$	e_6	e_7	$-e_8$
e_2	$3e_2$	0	$2e_1$	e_3	0	$-e_5$	e_8	0
e_3	$-2e_3$	$-e_1$	$-3e_4$	0	e_6	0	0	$-e_7$
e_4	0	0	0	0	0	0	0	0
e_5	$-2e_1$	$-3e_2$	$-e_3$	0	$-2e_5$	$-e_6$	$-e_7$	$-2e_8$
e_6	$2e_3$	e_1	$3e_4$	0	$-e_6$	0	0	e_7
e_7	$2e_3$	e_1	$3e_4$	0	$-e_6$	0	0	e_7
e_8	0	e_2	$-e_3$	$-2e_4$	0	$-e_6$	$-e_7$	0

The subalgebra spanned by the elements e_1, \dots, e_6 is the conservative (and, moreover, terminal) algebra W_2 of commutative 2-dimensional algebras. The subalgebra spanned by the elements e_1, \dots, e_4 is the conservative (and, moreover, terminal) algebra S_2 of all commutative 2-dimensional algebras with trace zero multiplication [20].

We now investigate the structure of $W(2)$ over fields of arbitrary characteristic. Regardless of $\text{char}(\mathbb{F})$, the diagram of $W(2)$ in the basis of the e_i 's is:



which is transitive (take into account that $-e_3^2 + ((e_3e_2)e_8)e_4 = e_4$). Now we claim:

Theorem 23. *For $\text{char}(\mathbb{F}) \neq 2, 3$ we have $\mathcal{M}(W(2)) = \text{End}_{\mathbb{F}}(W(2))$, hence $W(2)$ is simple. In the characteristic 2 case, $W(2)$ has a two dimensional ideal $I = \mathbb{F}(e_5 + e_8) \oplus \mathbb{F}(e_6 + e_7)$ and $W(2)/I \cong W_2$. Moreover $\mathcal{M}(W(2)) \cong M_6(\mathbb{F}) \oplus M_2(\mathbb{F})$. In the characteristic 3 case $\mathcal{M} = \mathcal{M}(W(2))$ has a 12-dimensional radical \mathcal{M}^\perp of square zero and $\mathcal{M}^\perp W(2) = \mathbb{F}(e_1 + e_5) \oplus \mathbb{F}(e_3 + e_6)$ is an ideal of $W(2)$.*

Proof. (A) Assume first that $\text{char}(\mathbb{F}) \neq 2, 3$. Then we have:

$$\begin{aligned} L_{e_8}L_{e_7}L_{e_8} &= 6E_{34}, \text{ implying } E_{3k} \in \mathcal{M} \text{ for any } k. \\ R_{e_7}^2 &= -E_{27}, \text{ implying } E_{2k} \in \mathcal{M} \text{ for any } k. \\ L_{e_7}^2 &= 6E_{14} + 2E_{23}, \text{ implying } E_{1k} \in \mathcal{M} \text{ for any } k. \\ L_{e_2}^2 &= 6E_{32} + 2E_{41}, \text{ implying } E_{4k} \in \mathcal{M} \text{ for any } k. \\ L_{e_2}R_{e_4} &= 3E_{13} + 2E_{21} - 2E_{83}, \text{ implying } E_{8k} \in \mathcal{M} \text{ for any } k. \\ L_{e_2}R_{e_6} &= -E_{15} + E_{55} + E_{85}, \text{ implying } E_{5k} \in \mathcal{M} \text{ for any } k. \\ L_{e_2} &= 3E_{12} + 2E_{31} + E_{43} - E_{65} + E_{78}, \text{ hence } -E_{65} + E_{78} \in \mathcal{M}. \end{aligned}$$

Thus $E_{7k} = (-E_{65} + E_{78})E_{8k} \in \mathcal{M}$ for any k . Therefore $E_{65} \in \mathcal{M}$ and so every $E_{6k} \in \mathcal{M}$. Thus we conclude $\mathcal{M}(W(2)) = \text{End}_{\mathbb{F}}(W(2))$ in the case of characteristic $\neq 2, 3$.

(B) Assume now $\text{char}(\mathbb{F}) = 2$. A simple but tedious computation reveals that the radical $R = \text{rad}(\langle \cdot, \cdot \rangle)$ of the trace form $\langle \cdot, \cdot \rangle : W(2) \times W(2) \rightarrow \mathbb{F}$ given as before by $\langle x, y \rangle := \text{trace}(xy)$ has a basis given by

$$\begin{aligned} r_1 &= E_{15} + E_{18}, & r_7 &= E_{45} + E_{48}, \\ r_2 &= E_{16} + E_{17}, & r_8 &= E_{46} + E_{47}, \\ r_3 &= E_{25} + E_{28}, & r_9 &= E_{55} + E_{58} + E_{85} + E_{88}, \\ r_4 &= E_{26} + E_{27}, & r_{10} &= E_{56} + E_{57} + E_{86} + E_{87}, \\ r_5 &= E_{35} + E_{38}, & r_{11} &= E_{65} + E_{68} + E_{75} + E_{78}, \\ r_6 &= E_{36} + E_{37}, & r_{12} &= E_{66} + E_{67} + E_{76} + E_{77}. \end{aligned}$$

It is also straightforward that $R^2 = 0$. The natural action $\mathcal{M} \times W(2) \rightarrow W(2)$ provides the two-dimensional ideal $R \cdot W(2) \triangleleft W(2)$ which is $R \cdot W(2) = \mathbb{F}(e_5 + e_8) \oplus \mathbb{F}(e_6 + e_7)$. Furthermore the quotient algebra $W(2)/RW(2)$ is isomorphic to the six-dimensional algebra B of section 2.2. By Theorem 12, $W(2)/RW(2)$ is simple. One can easily check that $RW(2) \subset \text{Lann}(W(2))$ but $RW(2) \not\subset \text{Rann}(W(2))$. On the other hand, the two-sided annihilator of the

ideal $RW(2)$, that is, the vector space of elements $x \in W(2)$ such that $x(RW(2)) = 0 = (RW(2))x$ is generated by $e_3 + e_7$, e_4 , $e_5 + e_8$ and $e_6 + e_7$. This implies that there is no ideal I complementing $RW(2)$ (because if I existed it would have dimension 6 and it would be contained in the linear span of $\{e_3 + e_7, e_4, e_5 + e_8, e_6 + e_7\}$ which is impossible). Now the natural representation $\mathcal{M} \rightarrow \text{End}(W(2))$ induces the isomorphism map

$$\begin{aligned} \mathcal{M} &\rightarrow \text{End}[W(2)/RW(2)] \times \text{End}(RW(2)) \\ T &\mapsto (\bar{T}, T|_{RW(2)}) \end{aligned}$$

where \bar{T} is the map induced in the quotient $W(2)/RW(2)$ by the fact that $RW(2)$ is \mathcal{M} -invariant. Hence, $\mathcal{M}(W(2)) \cong M_6(\mathbb{F}) \oplus M_2(\mathbb{F})$.

(C) In case $\text{char}(\mathbb{F}) = 3$ we have

$$\begin{aligned} L_{e_1} &= -E_{11} + E_{33} - E_{55} + E_{66} + E_{77} - E_{88}, \\ L_{e_2} &= -E_{31} + E_{43} - E_{65} + E_{78}, \\ L_{e_3} &= E_{13} - E_{21} + E_{56} - E_{87}, \\ L_{e_4} &= 0, \\ L_{e_5} &= E_{11} - E_{33} + E_{55} - E_{66} - E_{77} + E_{88}, \\ L_{e_6} &= -E_{13} + E_{21} - E_{56} + E_{87}, \\ L_{e_7} &= L_{e_6}, \\ L_{e_8} &= E_{22} - E_{33} + E_{44} - E_{66} - E_{77}, \\ R_{e_1} &= -E_{11} + E_{33} + E_{51} - E_{63} - E_{73}, \\ R_{e_2} &= -E_{31} + E_{61} + E_{71} + E_{82}, \\ R_{e_3} &= E_{13} - E_{21} - E_{53} - E_{83}, \\ R_{e_4} &= E_{23} + E_{84}, \\ R_{e_5} &= -E_{15} + E_{36} + E_{55} - E_{66} - E_{76}, \\ R_{e_6} &= E_{16} - E_{25} - E_{56} - E_{86}, \\ R_{e_7} &= E_{17} + E_{28} - E_{57} - E_{87}, \\ R_{e_8} &= -E_{18} - E_{37} + E_{58} + E_{67} + E_{77}. \end{aligned}$$

A basis for \mathcal{M} is given by the set of matrices:

$$\begin{aligned} &E_{ij} \text{ for } i = 2, 4, 7, 8 \text{ and any } j, \\ &E_{ai} + E_{bj} \text{ for } (a, b) \in \{(1, 5), (3, 6), (5, 5), (6, 6)\} \text{ and } (i, j) \in \{(1, 5), (3, 6)\}, \\ &E_{ai} - E_{bi} \text{ for } (a, b) \in \{(1, 5), (3, 6)\} \text{ and } i = 2, 4, 5, 6, 7, 8. \end{aligned}$$

We have computed again the radical \mathcal{M}^\perp of its trace form $\langle f, g \rangle := \text{Tr}(fg)$ and it is 12-dimensional. A basis for \mathcal{M}^\perp is

$$\begin{aligned} &E_{2,1} + E_{2,5}, \quad E_{2,3} + E_{2,6}, \quad E_{1,1} + E_{1,5} - E_{5,1} + 2E_{5,5}, \\ &E_{4,1} + E_{4,5}, \quad E_{4,3} + E_{4,6}, \quad E_{1,3} + E_{1,6} - E_{5,3} + 2E_{5,6}, \\ &E_{7,1} + E_{7,5}, \quad E_{7,3} + E_{7,6}, \quad E_{3,1} + E_{3,5} - E_{6,1} + 2E_{6,5}, \\ &E_{8,1} + E_{8,5}, \quad E_{8,3} + E_{8,6}, \quad E_{3,3} + E_{3,6} - E_{6,3} + 2E_{6,6}. \end{aligned}$$

One can easily check that $(\mathcal{M}^\perp)^2 = 0$. Furthermore

$$\mathcal{M}^\perp \cdot W(2) = \mathbb{F}(e_1 + e_5) \oplus \mathbb{F}(e_3 + e_6)$$

is a (two-sided) ideal of $W(2)$.

□

2.3.1. Automorphisms of $W(2)$. In this section we work again over a commutative ring R and denote $W(2)_R$ the R -algebra $\bigoplus_{i=1}^8 Re_i$ where the multiplication table of the e_i 's is that of the multiplication table of $W(2)$. If we take a generic element $w = \sum_{i=1}^8 \lambda_i e_i \in W(2)_R$ and compute the matrix of L_w relative to the basis of the e_i 's we obtain:

$$\begin{pmatrix} -\lambda_1 - 2\lambda_5 & 3\lambda_2 & 2\xi_1 & 0 & 0 & 0 & 0 & 0 & 0 \\ \xi_1 & -3\lambda_1 - 3\lambda_5 + \lambda_8 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 2\lambda_2 & 0 & \xi_2 & 3\xi_1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & \lambda_2 & 3\lambda_1 - 2\lambda_8 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -\lambda_1 - 2\lambda_5 & -\xi_1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -\lambda_2 & \xi_2 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & \xi_2 & \lambda_2 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & \xi_1 & -\lambda_1 - 2\lambda_5 & 0 \end{pmatrix},$$

where $\xi_1 = -\lambda_3 + \lambda_6 + \lambda_7$ and $\xi_2 = \lambda_1 - \lambda_5 - \lambda_8$, whose characteristic polynomial is invariant under automorphism so that L_w and $L_{\theta(w)}$ have the same characteristic polynomial for any automorphism θ of $W(2)_R$. We list some of the coefficients of that characteristic polynomial:

$$\begin{aligned} \Lambda_1(w) &:= 4(3\lambda_5 + \lambda_8), \\ \Lambda_2(w) &:= -4 \begin{pmatrix} 3\lambda_1^2 + 3\lambda_5\lambda_1 - 3\lambda_8\lambda_1 - 15\lambda_5^2 - \lambda_8^2 - 3\lambda_2\lambda_3 + 3\lambda_2\lambda_6 + \\ 3\lambda_2\lambda_7 - 12\lambda_5\lambda_8 \end{pmatrix}, \\ \Lambda_3(w) &:= -2(3\lambda_5 + \lambda_8) \begin{pmatrix} 18\lambda_1^2 + 18\lambda_5\lambda_1 - 18\lambda_8\lambda_1 - 27\lambda_5^2 + \lambda_8^2 - \\ 18\lambda_2\lambda_3 + 18\lambda_2\lambda_6 + 18\lambda_2\lambda_7 - 30\lambda_5\lambda_8 \end{pmatrix}. \end{aligned}$$

The left annihilator of $W(2)_R$ is

$$Re_4 \oplus R(2e_1 - e_5 + 3e_8) \oplus R(e_3 + e_6) \oplus R(e_3 + e_7).$$

Lemma 24. *Assume $W \neq 0$ to be a free R -module $W = \bigoplus_{i=1}^n Re_i$ and M a submodule with a basis $\{u_1, \dots, u_k\}$ which is a subset of another basis $\{u_1, \dots, u_n\}$ of W . Then if $M \subset \bigoplus_{i=1}^n \mathfrak{m}_i e_i$ for some maximal ideals $\mathfrak{m}_i \triangleleft R$ we have $M = 0$.*

Proof. For $i = 1, \dots, n$ define the R -algebras $K_i := R/\mathfrak{m}_i$ (which are fields) and $S = \bigotimes_{i=1}^n K_i$. If $M \neq 0$ the S -module $M \otimes_R S$ is free with a basis of cardinal k but for any $z \in M$ we have $z = \sum_{i=1}^n m_i e_i$ with $m_i \in \mathfrak{m}_i$ and any element $z \otimes 1_S \in M \otimes_R S$ satisfies

$$z \otimes 1_S = \sum_{i=1}^n m_i e_i \otimes 1_S = \sum_{i=1}^n e_i \otimes m_i 1_S$$

but

$$\begin{aligned} m_i 1_S &= m_i(1_1 \otimes \dots \otimes 1_n) = (1_1 \otimes \dots \otimes m_i 1_i \otimes \dots \otimes 1_n) = \\ &= (1_1 \otimes \dots \otimes 0 \otimes \dots \otimes 1_n) = 0. \end{aligned}$$

□

The fact the \mathfrak{m}_i is maximal in Lemma 24 is not important. What it is essential is that it is proper (as any maximal ideal is). So we could replace the maximality hypothesis in the Lemma with $\mathfrak{m}_i \neq R$.

It is also easily seen that if L is a free R -module with a (finite) basis $\{l_1, \dots, l_n\}$ then it may not have a system of generators of cardinal $< n$. This allows to extend the previous Lemma in the following sense:

Lemma 25. *Assume $W \neq 0$ to be a free R -module $W = \bigoplus_{i=1}^n Re_i$ and M a submodule with a basis $\{u_1, \dots, u_k\}$ which is a subset of another basis $\{u_1, \dots, u_n\}$ of W . Denote by $p_i: W \rightarrow R$ the i -th coordinate projection relative to the basis $\{e_i\}_{i=1}^n$. Then for each $i = 1, \dots, n$ if $p_i(R) \neq 0$ we have $p_i(W) = R$.*

Proof. Assume without loss of generality that the ideal $p_1(W)$ is proper and nontrivial. We can define the ring

$$S = R/p_1(W) \otimes \overbrace{R \otimes \dots \otimes R}^{n-1}$$

and $M_S := M \otimes S$ is a free S -module of dimension K but for any element $z \in M$ given by $z = \sum_{i=1}^n r_i e_i$ we have

$$z \otimes 1_S = r_1 e_1 \otimes 1_S + \sum_{i>1} r_i e_i \otimes 1_S = \sum_{i>1} e_i \otimes r_i$$

so that $\{e_i \otimes 1_S\}_{i>1}$ is a system of generators of M_S of cardinal $< n$. \square

It is known that any commutative ring R satisfies the strong rank condition [24, (1.38) Corollary, p. 15], equivalently, for any monomorphism $R^m \rightarrow R^n$ we have $m \leq n$. In particular consider the free R -module R^n with canonical basis $\{e_i\}_{i=1}^n$. If a free R -submodule M of R^n has $\dim_R(M) = k$ then $k < n$. Moreover if $\{e_1, \dots, e_k\} \subset M$ then we want also to prove that

$$M = \bigoplus_{i=1}^k Re_i. \tag{12}$$

Indeed, take a basis $\{u_i\}_{i=1}^k$ of M . Then for $1 \leq i \leq k$ we have $e_i = \sum_{q=1}^k a_i^q u_q$ and for any q we also have $u_q = \sum_{j=1}^n b_q^j e_j$ (where $a_i^q, b_q^j \in R$). Thus $1 = \sum_{q=1}^k a_i^q b_q^j = \delta_i^j$ (Kronecker's delta) or equivalently $AB = 1_k$ (identity matrix $k \times k$ in $M_n(R)$) where $A = (a_i^j)_{i,j=1}^k$ and $B = (b_i^j)_{i,j=1}^k$. But since R is a commutative ring, it is stably finite (see [24, (1.12) Proposition] and definition [24, §1B, p.5]). So $BA = 1_K$ also. Now denoting $\mathbf{u} := (u_1, \dots, u_k)$ and $\mathbf{e} := (e_1, \dots, e_k)$ we can write $A\mathbf{u}^t = \mathbf{e}^t$ hence $\mathbf{u}^t = B\mathbf{e}^t$ proving formula (12).

Lemma 26. *Assume $\frac{1}{2}, \frac{1}{3} \in R$ and that I is left ideal of $W(2)_R$ which is a free R -submodule and $\dim_R(I) = 4$. Denote by $p_i: W(2)_R \rightarrow R$ the i th coordinate function relative to the basis $\{e_i\}$. If $p_i(x) \in R^\times$ for some $x \in I$ and $i \in \{1, 2, 3, 4\}$ then $I = \bigoplus_{j=1}^4 Re_j$.*

Proof. First we prove that if some $e_i \in I$ (with $i \in \{1, 2, 3, 4\}$) then $I = \bigoplus_{j=1}^4 Rj$. Assume first that $e_1 \in I$, then in the first column of the table of

multiplication of $W(2)$ we can see that $e_2, e_3 \in I$ and since e_4 appears in third column we conclude $e_4 \in I$. So $\bigoplus_{i=1}^4 Re_i \subset I$ and $\dim_R(I) = 4$ implies by formula (12) that $I = \bigoplus_{i=1}^4 Re_i$. In case $e_2 \in I$ we can see that $e_1 \in I$ (second column of the table of multiplication of $W(2)$). The same applies if $e_3 \in I$. Finally if $e_4 \in I$ then $e_3 \in I$ for a similar reason. Now assume that some $x \in I$ has $p_1(x) \in R^\times$. We can assume without loss of generality that $p_1(x) = 1$. Since we have $e_8(e_2(e_5(e_1x))) = 6e_2$ then $e_2 \in I$ and we apply the proved part of the Lemma. If some $x \in I$ has $p_2(x) \in R^\times$ again we can assume $p_2(x) = 1$ and then since we have $e_3(e_3(e_3(e_1x))) = 18e_4$ we conclude $e_4 \in I$ and can apply again the proved part of the Lemma. In case $p_3(x) \in R^\times$ for some $x \in I$ we take into account that $e_8(e_2(e_2x)) = 6e_2$ implying $e_2 \in I$ (as before assuming $p_3(x) = 1$). Finally if $p_4(x) \in R^\times$ for some $x \in I$ we use $e_2(e_2(e_2x)) = 6e_2$. \square

Corollary 27. *Assume $\frac{1}{2}, \frac{1}{3} \in R$ and that I is left ideal of $W(2)_R$ which is a free R -submodule and $\dim_R(I) = 4$. Denote as before by $p_i: W(2)_R \rightarrow R$ the i th coordinate function relative to the basis $\{e_i\}_{i=1}^8$. The either $I = \bigoplus_{j=1}^4 Re_j$ or $I = \bigoplus_{j=5}^8 Re_j$.*

Proof. By Lemma 26 either $I = \bigoplus_{i=1}^4 Re_i$ or $I \subset \bigoplus_{i=5}^8 Re_i$. But $\dim_R(I) = 4$ so (12) gives the equality $I = \bigoplus_{i=5}^8 Re_i$. \square

Next we keep on assuming $\frac{1}{2}, \frac{1}{3} \in R$. We want to investigate the case that $\theta: W(2)_R \rightarrow W(2)_R$ be an automorphism such that

$$\theta(\bigoplus_{i=1}^4 Re_i) = \bigoplus_{i=5}^8 Re_i.$$

Denote $\theta(e_1) = \sum_{i=5}^8 \lambda_i e_i$. Since $\Lambda_1(e_1) = 0$ we have

$$\Lambda_1(\theta(e_1)) = 4(3\lambda_5 + \lambda_8) = 0 \text{ so } \lambda_8 = -3\lambda_5.$$

Also $\Lambda_2(e_1) = 3$ and $\Lambda_2(\theta(e_1)) = 12\lambda_5^2$ which implies $\lambda_5^2 = 1/4$ and in particular $\lambda_5 \in R^\times$. Furthermore $e_1^2 + e_1 = 0$ hence $\theta(e_1)^2 + \theta(e_1) = 0$ which (after the corresponding computation) gives $\lambda_5 = \frac{1}{2}$ and $\lambda_7 = 3\lambda_6$. Thus we have

$$\theta(e_1) = \frac{1}{2}e_5 + \lambda_6 e_6 + 3\lambda_6 e_7 - \frac{3}{2}e_8. \tag{13}$$

Next we study $\theta(e_2) = \sum_{i=5}^8 \mu_i e_i$. Again

$$\Lambda_1(e_2) = 0 = \Lambda_1(\theta(e_2)) = 4(3\mu_5 + \mu_8) \text{ hence } \mu_8 = -3\mu_5.$$

Moreover $\Lambda_2(e_2) = 0 = \Lambda_2(\theta(e_2)) = 12\mu_5^2$ hence $\mu_5^2 = \mu_8^2 = \mu_5\mu_8 = 0$. Since $e_2^2 = 0$ we have $0 = \theta(e_2)^2 = (\mu_6 - \mu_7)\mu_5 e_6 - (3\mu_6 + \mu_7)\mu_5 e_7$. Thus

$$\mu_6\mu_5 = \mu_7\mu_5 \text{ and } 3\mu_6\mu_5 = \mu_7\mu_5 \text{ whence } \mu_6\mu_5 = 0 = \mu_7\mu_5.$$

But then $\mu_5\theta(e_2) = 0$, that is, $\theta(\mu_5 e_2) = 0$ which gives $\mu_5 = 0$. We get

$$\theta(e_2) = \mu_6 e_6 + \mu_7 e_7. \tag{14}$$

But then, since $e_1 e_2 + 3e_2 = 0$, applying θ and taking into account (13) and (14), we find $0 = \theta(e_1)\theta(e_2) + 3\theta(e_2) = 4\mu_6 e_6 + 4\mu_7 e_7$ so that $\mu_6 = \mu_7 = 0$ which is a contradiction. So far we have proved that no automorphism of

$W(2)_R$ maps $\bigoplus_{i=1}^4 Re_i$ to $\bigoplus_{i=5}^8 Re_i$. As a consequence no automorphism of $W(2)_R$ maps $\bigoplus_{i=5}^8 Re_i$ to $\bigoplus_{i=1}^4 Re_i$.

Corollary 28. *If $\frac{1}{2}, \frac{1}{3} \in R$ any automorphism of $W(2)_R$ maps $\bigoplus_{i=1}^4 Re_i$ to itself and the same holds for $\bigoplus_{i=5}^8 Re_i$.*

Lemma 29. *Assume $\frac{1}{2}, \frac{1}{3} \in R$ and that I is left ideal of $W(2)_R$ which is a free R -submodule of dimension 2. Denote by $p_i: W(2)_R \rightarrow R$ the i th coordinate function relative to the basis $\{e_i\}$. Then $p_i(x) = 0$ for any $x \in I$ and $i \in \{1, 2, 3, 4\}$.*

Proof. First we assume that some $re_i \in I$ with $r \neq 0$ and $i \in \{1, 2, 3, 4\}$. This will take us to a contradiction. Indeed, under that assumption we have $I_r \supset R_r \frac{e_i}{1}$ where we denote by R_r the localization RS^{-1} being $S = \{1, r, r^2, \dots\}$. Let $W := W(2)_R$ and consider the localization $W_r := W \otimes_R R_r$ then (since R_r is a flat R -algebra) $I_r := I \otimes_R R_r$ is an ideal of W_r which a free R_r -module and $\dim_{R_r}(I_r) = 2$. We will identify W_r with the algebra of fractions $\frac{x}{r^n}$ ($x \in W, n \geq 0$) where $\frac{x}{r^n} = \frac{x'}{r^m}$ if and only if $r^k(r^m x - r^n x') = 0$ for some k . Now if $re_i \in I$ ($i = 1, 2, 3, 4$) then $\frac{re_i}{1} \in I_r$ so that $\frac{e_i}{1} \in I_r$. Consequently $R_r e_i \subset I_r$ and the multiplication table of W gives $\bigoplus_{i=1}^4 R_r e_i \subset I_r$ (we have identified $\frac{e_i}{1}$ with e_i). But then $4 \leq 2$ taking dimensions. We conclude that if $re_i \in I$ with $i \in \{1, 2, 3, 4\}$ then $r = 0$. Now consider $x \in I$ with $p_i(x) \neq 0$ and $i \in \{1, 2, 3, 4\}$. We have $I \ni e_8(e_2(e_5(e_1x))) = 6p_1(x)e_2$ whence $p_1(x) = 0$. Next we have $I \ni e_3(e_3(e_3(e_1x))) = 18p_2(x)e_4$ hence $p_2(x) = 0$. Then $I \ni e_8(e_2(e_2x)) = 6p_3(x)e_2$ whence $p_3(x) = 0$ and finally the equality $I \ni e_2(e_2(e_2x)) = 6p_4(x)e_2$ to deduce that $p_4(x) = 0$. \square

Consider now an automorphism θ of $W(2)_R$ (again $\frac{1}{2}, \frac{1}{3} \in R$) and let us study the image $\theta(e_6)$. Since $Re_5 \oplus Re_6$ is a left ideal of $W(2)_R$ and it is under the hypothesis of Lemma 29, we have $\theta(e_5), \theta(e_6) \in \bigoplus_{i=5}^8 Re_i$. So for instance $\theta(e_5) = \sum_{i=5}^8 \mu_i e_i$ and $\theta(e_6) = \sum_{i=5}^8 \lambda_i e_i$ and we can use again the invariants Λ_1 and Λ_2 . We have

$$12 = \Lambda_1(e_5) = \Lambda_1(\theta(e_5)) = 4(3\mu_5 + \mu_8)$$

whence $\mu_8 = 3 - 3\mu_5$. Also $-15 = \Lambda_2(e_5) = \Lambda_2(\theta(e_5)) = 12\mu_5^2 - 18\mu_5 - 9$. So we deduce that $2\mu_5^2 - 3\mu_5 + 1 = 0$ implying that μ_5 is invertible. Furthermore, $\theta(e_5)^2 + 2\theta(e_5) = 0$ hence the following elements of R are zero:

$$2(\mu_5 - 1)\mu_5, \mu_5\mu_6 - \mu_6 - \mu_5\mu_7, -3\mu_5\mu_6 + 3\mu_6 - \mu_5\mu_7 + 2\mu_7, 6(\mu_5 - 1)^2.$$

This implies that $\mu_5 = 1, \mu_7 = 0$ and $\mu_8 = 0$. So $\theta(e_5) = e_5 + \mu_6 e_6$. Now using again the invariants $\Lambda_1(e_6) = 0$ and $\Lambda_2(e_6) = 0$. We have

$$\Lambda_1(\theta(e_6)) = 4(3\lambda_5 + \lambda_8) = 0 \text{ whence } \lambda_8 = -3\lambda_5.$$

Also $\Lambda_2(\theta(e_6)) = 0$ from which we derive $\lambda_5^2 = 0$ and consequently $\lambda_8^2 = 0 = \lambda_8\lambda_5$. We also have $\theta(e_6)^2 = 0$ which gives

$$\lambda_5\lambda_6 - \lambda_5\lambda_7 = 0, -3\lambda_5\lambda_6 - \lambda_5\lambda_7 = 0 \text{ and so } \lambda_5\lambda_6 = \lambda_5\lambda_7 = 0.$$

As a consequence $\lambda_5\theta(e_6) = 0$ which gives $\lambda_5 = 0$. So $\theta(e_6) = \lambda_6e_6$ and in summary we have

$$\begin{cases} \theta(e_5) = e_5 + \mu_6e_6 \\ \theta(e_6) = \lambda_6e_6. \end{cases} \tag{15}$$

Now, under the same assumptions $\frac{1}{2}, \frac{1}{3} \in R$ let us investigate $\theta(e_7), \theta(e_8)$ for $\theta \in \text{aut}(W(2)_R)$. As in the previous case we have $\theta(e_7), \theta(e_8) \in \oplus_{i=5}^8 Re_i$. Write $\theta(e_7) = \sum_{i=5}^8 \gamma_i e_i$, since $\theta(e_6)\theta(e_7) = 0$ we get $\gamma_5 = \gamma_8 = 0$ so that $\theta(e_7) = \gamma_6e_6 + \gamma_7e_7$. Finally write $\theta(e_8) = \sum_{i=5}^8 \delta_i e_i$, from the equality $\Lambda_1(e_8) = \Lambda_1(\theta(e_8))$ we get $\delta_8 = 1 - 3\delta_5$ and from $\Lambda_2(e_8) = \Lambda_2(\theta(e_8))$ we have $\delta_5(2\delta_5 - 1) = 0$. Now the couple of identities $e_6e_8 = e_7$ and $e_8e_6 = -e_6$ give the equations

$$-\gamma_6 - \delta_5\lambda_6 = 0, \quad -\gamma_7 - 3\delta_5\lambda_6 + \lambda_6 = 0, \quad 2\delta_5\lambda_6 = 0$$

so that $\gamma_6 = 0, \gamma_7 = \lambda_6$ and $\delta_5 = 0$ (because $\lambda_6 \neq 0$). Then $\theta(e_7) = \lambda_6e_7$ and $\theta(e_8) = \delta_6e_6 + \delta_7e_7 + e_8$ but since $\theta(e_8)^2 = 0$ we get $\delta_6 = 0$ so that

$$\begin{cases} \theta(e_7) = \lambda_6e_7 \\ \theta(e_8) = \delta_7e_7 + e_8. \end{cases} \tag{16}$$

Thus we conclude

Proposition 30. *If $\frac{1}{2}, \frac{1}{3} \in R$ any automorphism θ of $W(2)_R$ fixes any of the left ideals $\oplus_{i=1}^4 Re_i, \oplus_{i=1}^6 Re_i, \oplus_{i=5}^6 Re_i$ and $\oplus_{i=7}^8 Re_i$. Relative to the basis $\{e_i\}_{i=1}^8$ the matrix of an automorphism is of the form*

$$\left(\begin{array}{cccccc|cc} 1 & 0 & 2tx & 3t^2x^2 & 0 & 0 & 0 & 0 \\ x & \frac{1}{t} & tx^2 & t^2x^3 & 0 & 0 & 0 & 0 \\ 0 & 0 & t & 3t^2x & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & t^2 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & -tx & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & t & 0 & 0 \\ \hline 0 & 0 & 0 & 0 & 0 & 0 & t & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & tx & 1 \end{array} \right) \tag{17}$$

Proof. Since θ restricts to an automorphism of $W_2(R) = \oplus_{i=1}^6 Re_i$, the 6×6 upper left block in (17) is in (8). It remains to prove that $\theta(e_8) = tx e_7 + e_8$ but we have proved in (16) that $\theta(e_8) = \delta_7 e_7 + e_8$. Since $\theta(e_8)\theta(e_3) + \theta(e_3) = 0$ we have

$$0 = (\delta_7 e_7 + e_8)(te_3 + 3t^2xe_4) + te_3 + 3t^2xe_4 = 3\delta_7te_4 - te_3 - 6t^2xe_4 + te_3 + 3t^2xe_4 = 3\delta_7te_4 - 3t^2xe_4$$

and since t is invertible $\delta_7 = tx$. □

2.3.2. The case $2R = 0$. Note that necessarily $\frac{1}{3} \in R$. In this case define $f_1 := e_5 + e_8$ and $f_2 = e_6 + e_7$. Then $I := Rf_1 \oplus Rf_2$ is a 2-dimensional (two-sided) ideal of $W(2)_R$ (see Theorem 23). It has a basis $\{f_1, f_2\}$ which is a subbasis of $\{e_1, e_2, e_3, e_4, e_5, e_6, e_6 + e_7, e_5 + e_8\}$ which can be seen to be a basis of $W(2)_R$. Indeed the matrix of coordinates of these vectors relative to the basis of the $\{e_i\}$ is $\begin{pmatrix} I_6 & 0 \\ M & I_2 \end{pmatrix}$ where I_6 and I_2 denote the identity matrices of size 6 and 2 respectively and $M = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 1 & 0 \end{pmatrix}$. It is easy to check that M is invertible and agrees with its own inverse. The ideal I satisfies $IW(2)_R = 0$.

Lemma 31. *Assume that $2R = 0$ and $J \triangleleft W(2)_R$ is a 2-dimensional ideal such that $JW(2)_R = 0$ then $J \subset I$.*

Proof. Any $x \in J$ satisfies $xW(2)_R = 0$ which implies that the elements of J are of the form

$$g = \lambda_3 e_3 + \lambda_4 e_4 + \lambda_5 (e_5 + e_8) + \lambda_6 e_6 + (\lambda_3 + \lambda_6) e_7.$$

Note that the 5th and 8th coordinates of g (relative to the basis $\{e_i\}$) agree. So since $e_2 g \in J$ we must have $\lambda_6 = p_5(e_2 g) = p_8(e_2 g) = \lambda_3 + \lambda_6$ whence $\lambda_3 = 0$ and we have proved that the elements of J satisfy $p_3(J) = 0$. So a general element of J is of the form $g = \lambda_4 e_4 + \lambda_5 (e_5 + e_8) + \lambda_6 (e_6 + e_7)$. But $J \ni e_2 g = \lambda_4 e_3 + \lambda_6 e_5 + (\lambda_3 + \lambda_6) e_8$ which implies $\lambda_4 = 0$. Thus $g \in R(e_5 + e_8) \oplus R(e_6 + e_7)$. □

Under the hypothesis in the title of this subsection, if $\theta \in \text{aut}(W(2)_R)$, Lemma 31 implies $\theta(I) \subset I$ (recall that I is the ideal $I = Rf_1 \oplus Rf_2$ defined above). Consequently $I \subset \theta^{-1}(I) \subset I$ so that $\theta(I) = I$ for any $\theta \in \text{aut}(W(2)_R)$. Since $W(2)_R/I \cong W_2$ (see Theorem 23) any $\theta \in \text{aut}(W(2)_R)$ induces an automorphism $\bar{\theta}: W_2 \rightarrow W_2$. Then the matrix of $\bar{\theta}$ relative to the basis $\{\bar{e}_i\}_{i=1}^6$ (begin $\bar{e}_i := e_i + I$) is the one in formula (8). So the matrix of θ relative to the basis $\{e_1, \dots, e_6, f_1, f_2\}$ of $W(2)_R$ is of the form

$$\left(\begin{array}{cccccc|cc} 1 & 0 & 0 & t^2 x^2 & 0 & 0 & a_1 & a_2 \\ x & \frac{1}{t} & tx^2 & t^2 x^3 & 0 & 0 & a_3 & a_4 \\ 0 & 0 & t & t^2 x & 0 & 0 & a_5 & a_6 \\ 0 & 0 & 0 & t^2 & 0 & 0 & a_7 & a_8 \\ 0 & 0 & 0 & 0 & 1 & tx & a_9 & a_{10} \\ 0 & 0 & 0 & 0 & 0 & t & a_{11} & a_{12} \\ \hline 0 & 0 & 0 & 0 & 0 & 0 & a_{13} & a_{14} \\ 0 & 0 & 0 & 0 & 0 & 0 & a_{15} & a_{16} \end{array} \right) \tag{18}$$

where $t \in R^\times$. Furthermore, if we write the matrix of θ relative to the basis of the e_i 's and impose the conditions for automorphism we find the relations

$$\begin{aligned} a_1 = a_2 = a_3 = a_4 = a_5 = a_6 = a_7 = a_8 = 0, a_9 = a_{10}tx, \\ a_{11} = a_{10}t, a_{12} = 0, a_{14} = 0, a_{15} = a_{13}x, a_{16} = \frac{a_{13}}{t}. \end{aligned}$$

Lemma 32. *In case $2R = 0$ the matrix of an automorphism $\theta \in \text{aut}(W(2)_R)$ relative to the basis $\{e_1, \dots, e_6, f_1, f_2\}$ of $W(2)_R$ is*

$$\Omega_{t,x,v,u} = \begin{pmatrix} 1 & 0 & 0 & t^2x^2 & 0 & 0 & 0 & 0 \\ x & \frac{1}{t} & tx^2 & t^2x^3 & 0 & 0 & 0 & 0 \\ 0 & 0 & t & t^2x & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & t^2 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & tx & utx & u \\ 0 & 0 & 0 & 0 & 0 & t & ut & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & v & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & vx & \frac{v}{t} \end{pmatrix} \quad (19)$$

where we have replaced a_{10} with u and a_{13} with v . Furthermore $t, v \in R^\times$, $x, u \in R$.

We have the relations

$$\Omega_{t,x,v,u}\Omega_{t',x',v',u'} = \Omega_{tt',x+x'/t,vv',u'+uv'/t'} \text{ and } \Omega_{t,x,v,u}^{-1} = \Omega_{1/t,tx,1/v,tu/v}.$$

The set $G_1 := \{\Omega_{t,x,1,0} : t \in R^\times, x \in R\}$ is a subgroup of $\text{aut}(W(2)_R)$ isomorphic to $\text{Aff}_2(R)$. Indeed, if we consider

$$\text{Aff}_2(R) = \left\{ \begin{pmatrix} 1 & x \\ 0 & t \end{pmatrix} : t \in R^\times, x \in R \right\}$$

we have a group isomorphism $\gamma_1: \text{Aff}_2(R) \rightarrow G_1$ such that

$$\begin{pmatrix} 1 & x \\ 0 & t \end{pmatrix} \mapsto \Omega_{t,t^{-1}x,1,0}.$$

On the other hand $G_2 := \{\Omega_{1,0,v,u} : v \in R^\times, u \in R\}$ is also a subgroup of $\text{aut}(W(2)_R)$ isomorphic to $\text{Aff}_2(R)$ via the map $\gamma_2: \text{Aff}_2(R) \rightarrow G_2$ such that $\begin{pmatrix} 1 & x \\ 0 & t \end{pmatrix} \mapsto \Omega_{1,0,t,x}$. It is easily seen that G_2 is a normal subgroup of $\text{aut}(W(2)_R)$ and the map $\rho: G_1 \rightarrow \text{aut}(G_2)$ given by

$$\rho(\Omega_{t,x,1,0})(\Omega_{1,0,v,u}) = \Omega_{t,x,1,0}\Omega_{1,0,v,u}\Omega_{t,x,1,0}^{-1} = \Omega_{1,0,v,tu}$$

is a group homomorphism. We also have $\Omega_{t,x,v,u} = \Omega_{t,x,1,0}\Omega_{1,0,v,u}$ and so $\text{Aut}(W(2)_R) = G_2 \rtimes G_1$ with multiplication

$$(g_2g_1)(g'_2g'_1) = [g_2\rho(g_1)(g'_2)](g_1g'_1).$$

If we define $\tau_{x,t} := \begin{pmatrix} 1 & x \\ 0 & t \end{pmatrix}$ so that $\text{Aff}_2(R) = \{\tau_{x,t} : x \in R, t \in R^\times\}$ then we have an action of $\text{Aff}_2(R)$ on itself by automorphisms $\rho': \text{Aff}_2(R) \rightarrow \text{aut}(\text{Aff}_2(R))$ given by $\rho'(\tau_{x,t})(\tau_{u,v}) = \tau_{tu,v}$. Then there is a commutative square

$$\begin{array}{ccc} G_1 & \xrightarrow{\rho} & \text{aut}(G_2) \\ \gamma_1 \downarrow & & \downarrow \text{inn}_{\gamma_2} \\ \text{Aff}_2(R) & \xrightarrow{\rho'} & \text{aut}(\text{Aff}_2(R)) \end{array} \quad \begin{array}{c} \theta \\ \downarrow \\ \gamma_2^{-1}\theta\gamma_2 \end{array}$$

and we conclude that $\text{aut}(W(2)_R) \cong \text{Aff}_2(R) \rtimes \text{Aff}_2(R)$.

2.3.3. The case $3R = 0$. Note that necessarily $\frac{1}{2} \in R$. Consider an R -algebra A which is a free R -module with a finite basis. Let $\mathcal{M} := \mathcal{M}(A)$ be its multiplication algebra and $\text{Tr}: \mathcal{M} \rightarrow R$ the trace (so $\text{Tr}(T)$ is the trace of the matrix of T relative to any basis of the R -module A). Also denote by $k: \mathcal{M} \times \mathcal{M} \rightarrow R$ the symmetric R -bilinear map $k(T, S) := \text{Tr}(TS)$. This satisfies $k(TT', S) = k(T, T'S) = K(T', ST)$ for any $T, T', S \in \mathcal{M}$. Thus $\mathcal{M}^\perp := \{T \in \mathcal{M}: k(T, _) = 0\}$ is an ideal of \mathcal{M} and $\mathcal{M}^\perp A$ an ideal of A . There is also an action $\text{aut}(A) \times \mathcal{M} \rightarrow \mathcal{M}$ such that $\varphi \cdot T = T^* := \varphi T \varphi^{-1}$ for any $\varphi \in \text{aut}(A)$ and $T \in \mathcal{M}$. Furthermore $k(T^*, S^*) = K(T, S)$ for any $S, T \in \mathcal{M}$ so that $(\mathcal{M}^\perp)^* \subset \mathcal{M}^\perp$ or equivalently $\text{aut}(A) \cdot \mathcal{M}^\perp \subset \mathcal{M}^\perp$. Consequently the ideal $\mathcal{M}^\perp A$ of A is invariant under automorphisms of A : for any $\varphi \in \text{aut}(A)$, $T \in \mathcal{M}^\perp$ and $a \in A$ one has $\varphi(T(a)) = T^* \varphi(a) \in \mathcal{M}^\perp A$.

Remark 33. Let \mathbb{F} be an arbitrary field in this Lemma and U be a finite-dimensional \mathbb{F} -algebra, $\mathcal{M} = \mathcal{M}(U)$ its multiplication algebra, $I \triangleleft \mathcal{M}$ and $R \in \text{Alg}_{\mathbb{F}}$. If $j: IU \rightarrow U$ is the inclusion, identifying $IU \otimes R$ with $(IU)_R$ via $j \otimes 1_R: IU \otimes R \rightarrow U_R$, we have $(IU)_R = I_R U_R$.

We now particularize considering $W(2)_R$. We start with $W(2)$ over a field \mathbb{F} of characteristic 3 and take $A = W(2)_R$. If we denote $\mathcal{M} = \mathcal{M}(W(2))$ then \mathcal{M}_R can be identified with $\mathcal{M}(W(2)_R)$ ([6, (2.5) Lemma (a)]). Also we have $k: \mathcal{M} \times \mathcal{M} \rightarrow \mathbb{F}$ as above: $k(T, S) = \text{Tr}(TS)$ inducing $k_R: \mathcal{M}_R \times \mathcal{M}_R \rightarrow R$ and we have the standard result that $(\mathcal{M}_R)^\perp \cong (\mathcal{M}^\perp)_R$. By Theorem 23 we have $\mathcal{M}^\perp W(2) = \mathbb{F}(e_1 + e_5) \oplus \mathbb{F}(e_3 + e_6)$ hence by Remark 33,

$$\mathcal{M}_R^\perp W(2)_R = R(e_1 + e_5) \oplus R(e_3 + e_6).$$

So this ideal is invariant under automorphisms of $W(2)_R$.

Next we compute the quotient algebra $W(2)_R/I$ where

$$I = R(e_1 + e_5) \oplus R(e_3 + e_6).$$

We consider a basis of $W(2)_R/I$ given by

$$\begin{cases} f_i = e_i + I, \text{ for } i = 1, 2, 3, 4, \\ f_5 = e_8 + I, \\ f_6 = 2e_7 + I. \end{cases}$$

The multiplication of the quotient algebra relative to this basis is given in the following table

	f_1	f_2	f_3	f_4	f_5	f_6
f_1	$2f_1$	0	f_3	0	$2f_5$	f_6
f_2	0	0	$2f_1$	f_3	0	$2f_5$
f_3	f_3	$2f_1$	0	0	f_6	0
f_4	0	0	0	0	0	0
f_5	f_1	0	$2f_3$	0	f_5	$2f_6$
f_6	$2f_3$	f_1	0	0	$2e_6$	0

So we conclude that $W(2)_R/I \cong W_2(R)$ and any automorphism of $W(2)_R$ induces an automorphism of $W_2(R)$ whose matrix relative to the basis of the f_i 's is given in (10). Consequently any automorphism θ of $W(2)_R$ acts in the form

$$\begin{aligned} \theta(e_1) &= e_1 + ce_3 + (e_1 + e_5)t_1 + (e_3 + e_6)t_2 \\ &= (t_1 + 1)e_1 + (c + t_2)e_3 + t_1e_5 + t_2e_6, \\ \theta(e_2) &= -\frac{c}{a}e_1 + \frac{1}{a}e_2 + \frac{c^2}{a}e_3 - \frac{c^3}{a}e_4 + t_3(e_1 + e_5) + t_4(e_3 + e_6) \\ &= (t_3 - \frac{c}{a})e_1 + \frac{1}{a}e_2 + (\frac{c^2}{a} + t_4)e_3 - \frac{c^3}{a}e_4 + t_3e_5 + t_4e_6, \\ \theta(e_3) &= ae_3 + t_5(e_1 + e_5) + t_6(e_3 + e_6) \\ &= (a + t_6)e_3 + t_5e_1 + t_5e_5 + t_6e_6, \\ \theta(e_4) &= a^2e_4 + t_7(e_1 + e_5) + t_8(e_3 + e_6) \\ &= t_7e_1 + t_8e_3 + a^2e_4 + t_7e_5 + t_8e_6. \end{aligned}$$

But imposing the conditions $\theta(e_i e_j) = \theta(e_i)\theta(e_j)$ for $i, j \in \{1, 2, 3, 4\}$ we get

$$t_2 = ct_1, t_i = 0 \text{ for } i \geq 3.$$

Thus, the coordinates of $\theta(e_i)$, $i = 1, 2, 3, 4$ relative to the $\{e_j\}_{j=1}^8$ written in matrix form give

$$\begin{pmatrix} 1 & 0 & c & 0 & 0 & 0 & 0 & 0 \\ -\frac{c}{a} & \frac{1}{a} & \frac{c^2}{a} & -\frac{c^3}{a} & 0 & 0 & 0 & 0 \\ 0 & 0 & a & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & a^2 & 0 & 0 & 0 & 0 \end{pmatrix}.$$

On the other hand since the image of $e_1 + e_5$ and $e_3 + e_6$ is in $R(e_1 + e_5) \oplus R(e_3 + e_6)$ we have

$$\begin{aligned} \theta(e_5) &= -\theta(e_1) + x_1(e_1 + e_5) + x_2(e_3 + e_6) \\ &= (-c - t_2 + x_2)e_3 + (-t_1 + x_1 - 1)e_1 + (x_1 - t_1)e_5 + (x_2 - t_2)e_6, \\ \theta(e_6) &= -\theta(e_3) + x_3(e_1 + e_5) + x_4(e_3 + e_6) \\ &= (-a - t_6 + x_4)e_3 + (x_3 - t_5)e_1 + (x_3 - t_5)e_5 + (x_4 - t_6)e_6. \end{aligned}$$

Imposing the conditions $\theta(e_i e_j) = \theta(e_i)\theta(e_j)$ for $i \in \{1, 2, 3, 4\}$ and $j \in \{5, 6\}$ we get

$$t_1 = \frac{t_6}{a}, \quad x_2 = cx_1, \quad x_3 = 0, \quad x_4 = \frac{x_1}{a}$$

and the coordinates of $\theta(e_i)$ with $i = 1, \dots, 6$ relative to the basis $\{e_j\}_{j=1}^8$ written in a matrix form are

$$\begin{pmatrix} 1 & 0 & c & 0 & 0 & 0 & 0 & 0 \\ \frac{2c}{a} & \frac{1}{a} & \frac{c^2}{a} & \frac{2c^3}{a} & 0 & 0 & 0 & 0 \\ 0 & 0 & a & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & a^2 & 0 & 0 & 0 & 0 \\ b-1 & 0 & c(b-1) & 0 & b & bc & 0 & 0 \\ 0 & 0 & a(b-1) & 0 & 0 & ab & 0 & 0 \end{pmatrix}.$$

Finally writing $\theta(e_7) = \sum \lambda_i e_i$ and $\theta(e_8) = \sum \mu_i e_i$ and imposing the conditions $\theta(e_i e_j) = \theta(e_i)\theta(e_j)$ for $i \in \{1, \dots, 8\}$ and $j \in \{7, 8\}$ we get the matrix of a general automorphism $\theta \in \text{aut}_R(W(2)_R)$ which is

$$M_{a,b,c,k} := \begin{pmatrix} 1 & 0 & c & 0 & 0 & 0 & 0 & 0 \\ -\frac{c}{a} & \frac{1}{a} & \frac{c^2}{a} & -\frac{c^3}{a} & 0 & 0 & 0 & 0 \\ 0 & 0 & a & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & a^2 & 0 & 0 & 0 & 0 \\ b-1 & 0 & bc-c & 0 & b & bc & 0 & 0 \\ 0 & 0 & ab-a & 0 & 0 & ab & 0 & 0 \\ 0 & 0 & ak & 0 & 0 & ak & a & 0 \\ -k & 0 & -ck & 0 & -k & -ck & -c & 1 \end{pmatrix} \quad (20)$$

with $a, b \in R^\times$, $k, c \in R$.

Theorem 34. *If $3R = 0$ the matrix of any automorphism of $W(2)_R$ relative to a basis $\{e_i\}$ with multiplication table as in the table of multiplication of $W(2)$ is of the form (20) with $a, b \in R^\times$, $c, k \in R$.*

We have

$$M_{a,b,c,d} M_{a',b',c',d'} = M_{aa',bb',ca'+c',db'+d}, \quad M_{a,b,c,d}^{-1} = M_{\frac{1}{a}, \frac{1}{b}, -\frac{c}{a}, -\frac{k}{b}},$$

then $\text{aut}_R(W(2)_R) \cong \text{Aff}_2(R) \times \text{Aff}_2(R)$ via the isomorphism

$$M_{a,b,c,d} \mapsto \left(\begin{pmatrix} 1 & c \\ 0 & a \end{pmatrix}, \begin{pmatrix} 1 & d \\ 0 & b \end{pmatrix} \right)$$

and as an affine group scheme

$$\mathbf{aut}(W(2)) \cong \mathbf{Aff}_2 \times \mathbf{Aff}_2.$$

In this case, we have a direct product unlike the case $2R = 0$ in which the product was semidirect.

References

- [1] F. Arzikulov, N. Umrzaqov, *Conservative algebras of 2-dimensional algebras. III*, Communications in Mathematics, **29**:2 (2021), 255–267. Zbl 1477.17038
- [2] A. Calderón, A. Fernández Ouaridi, I. Kaygorodov, *The classification of 2-dimensional rigid algebras*, Linear and Multilinear Algebra, **68**:4 (2020), 828–844. Zbl 1471.17002
- [3] N. Cantarini, V. Kac, *Classification of linearly compact simple rigid superalgebras*, International Mathematics Research Notices, **17** (2010), 3341–3393. Zbl 1269.17002
- [4] R. Fehlberg Júnior, I. Kaygorodov, *On the Kantor product. II*, Carpathian Mathematical Publications, **14**:2 (2022), 543–563. Zbl 1528.17001
- [5] A. Fernández Ouaridi, I. Kaygorodov, C. Martín González, *Conservative algebras of 2-dimensional algebras. IV*, Journal of Algebra and its Applications, **24**:6 (2025), 2550143. Zbl 8011281
- [6] D. R. Finston, *On multiplication algebras*, Transactions of the American Mathematical Society, **293**:2 (1986), 807–818. Zbl 0609.16018
- [7] A. Giambruno, S. Mishchenko, M. Zaicev, *Codimension growth of two-dimensional non-associative algebras*, Proceedings of the American Mathematical Society, **135**:11 (2007), 3405–3415. Zbl 1156.17001

- [8] N. Jacobson, *A note on non-associative algebras*, Duke Mathematical Journal, **3** (1937), 544–548. Zbl 0018.05005
- [9] N. Jacobson, *Structure of Rings*, American Mathematical Society Colloquium Publications, Vol. 37. Revised edition American Mathematical Society, Providence, R.I. 1964 ix+299 pp.
- [10] I. Kantor, *Certain generalizations of Jordan algebras. (Russian)*, Proceedings of the Seminar on Vector and Tensor Analysis, **16** (1972), 407–499. Zbl 0272.17001
- [11] I. Kantor, *A universal attracting object in the category of conservative algebras. (Russian)*, Proceedings of the Seminar on Vector and Tensor Analysis, **23** (1988), 45–48. Zbl 0809.17030
- [12] I. Kantor, *On an extension of a class of Jordan algebras*, Algebra and Logic, **28:2** (1989), 117–121. Zbl 0704.17017
- [13] I. Kantor, *Terminal trilinear operations*, Algebra and Logic, **28** (1989), 1, 25–40. Zbl 0693.17003
- [14] I. Kantor, *Some problems in L-functor theory. (Russian)*, Proceedings of the Institute of Mathematics of the Siberian Branch of the USSR Academy of Sciences (Novosibirsk), vol. **16** (1989), Research on Ring and Algebra Theory, pp. 54–75.
- [15] I. Kantor, *An universal conservative algebra*, Siberian Mathematical Journal, **31:3** (1990), 388–395. Zbl 0711.17003
- [16] I. Kaygorodov, *On the Kantor Product*, Journal of Algebra and Its Applications, **16:9** (2017), 1750167.
- [17] Kaygorodov I., *Non-associative algebraic structures: classification and structure*, Communications in Mathematics, **32:3** (2024), 1–62. Zbl 1544.17001
- [18] I. Kaygorodov, M. Khrypchenko, Yu. Popov, *The algebraic and geometric classification of nilpotent terminal algebras*, Journal of Pure and Applied Algebra, **225:6** (2021), 106625. Zbl 1471.17005
- [19] I. Kaygorodov, A. Khudoyberdiyev, A. Sattarov, *One-generated nilpotent terminal algebras*, Communications in Algebra, **48:10** (2020), 4355–4390. Zbl 1471.17006
- [20] I. Kaygorodov, A. Lopatin, Yu. Popov, *Conservative algebras of 2-dimensional algebras*, Linear Algebra and its Applications, **486** (2015), 255–274. Zbl 1321.17002
- [21] I. Kaygorodov, Yu. Popov, A. Pozhidaev, *The universal conservative superalgebra*, Communications in Algebra, **47:10** (2019), 4066–4076. Zbl 1475.17001
- [22] I. Kaygorodov, Yu. Volkov, *Conservative algebras of 2-dimensional algebras. II*, Communications in Algebra, **45:8** (2017), 3413–3421. Zbl 1427.17003
- [23] I. Kaygorodov, Yu. Volkov, *The variety of 2-dimensional algebras over algebraically closed fields*, Canadian Journal of Mathematics, **71:4** (2019), 819–842. Zbl 1440.17027
- [24] T. Y. Lam, *Lectures on Modules and Rings*, Graduate Texts in Mathematics 189, Springer, 1999. Zbl 0911.16001
- [25] H. Petersson, *The classification of two-dimensional nonassociative algebras*, Results in Mathematics, **37:1-2** (2000), 120–154. Zbl 0956.17002
- [26] Yu. Popov, *Conservative algebras and superalgebras: a survey*, Communications in Mathematics, **28:2** (2020), 231–251. Zbl 1482.17001

IVAN KAYGORODOV

CMA-UBI, UNIVERSITY OF BEIRA INTERIOR, PORTUGAL.

MOSCOW CENTER FOR FUNDAMENTAL AND APPLIED MATHEMATICS, RUSSIA.

Email address: kaygorodov.ivan@gmail.com

DOLORES MARTIN BARQUERO

DEPARTAMENTO DE MATEMÁTICA APLICADA, ESCUELA DE INGENIERÍAS INDUSTRI-
ALES, UNIVERSIDAD DE MÁLAGA, CAMPUS DE TEATINOS, SPAIN.

Email address: dmartin@uma.es

CANDIDO MARTÍN GONZÁLEZ

DEPARTAMENTO DE MATEMÁTICA APLICADA, ESCUELA DE INGENIERÍAS INDUSTRI-
ALES, UNIVERSIDAD DE MÁLAGA, CAMPUS DE TEATINOS, SPAIN.

Email address: candido_m@uma.es